

**LIGHT SUBGRAPHS OF GRAPHS EMBEDDED IN
THE PLANE AND IN THE PROJECTIVE PLANE
– A SURVEY –**

S. JENDROL' ¹⁾ AND H.-J. VOSS ^{2)†}

¹⁾ Institute of Mathematics, Faculty of Science,
P.J.Šafárik University,
Jesenná 5, 041 54 Košice, Slovakia
jendrol@kosice.upjs.sk

²⁾ Department of Algebra, Technical University Dresden,
Mommensenstrasse 13. D-01062 Dresden, Germany
voss@math.tu-dresden.de

ABSTRACT. It is well known that every planar graph contains a vertex of degree at most 5. A beautiful theorem of Kotzig states that every 3-connected planar graph contains an edge whose endvertices have degree-sum at most 13. Recently, Fabrici and Jendroľ proved that every 3-connected planar graph G that contains a k -path, a path on k vertices, contains also a k -path P such that every vertex of P has degree at most $5k$. A beautiful result by Enomoto and Ota says that every 3-connected planar graph G of order at least k contains a connected subgraph H of order k such that the degree sum of vertices of H in G is at most $8k - 1$. Motivated by these results, a concept of light graphs has been introduced. A graph H is said to be *light* in a class \mathcal{G} of graphs if at least one member of \mathcal{G} contains a copy of H and there is an integer $w(H, \mathcal{G})$ such that each member G of \mathcal{G} with a copy of H also has a copy of H with degree sum $\sum_{v \in V(H)} \deg_G(v) \leq w(H, \mathcal{G})$.

In this paper we present a survey of results on light graphs in different families of plane and projective plane graphs and multigraphs. A similar survey dealing with the family of all graphs embedded in surfaces other than the plane and the projective plane has been prepared as well.

CONTENTS

1. Introduction	2
2. Notation and preliminaries	4
3. Light edges	6

Key words and phrases. 3-connected planar graphs, projective planar graphs, polyhedral maps, light subgraphs, path, cycle, embeddings, subgraphs with restricted degrees, triangulation, 2-manifolds
2000 Mathematical Subject Classification: Primary: 05C10 Secondary: 52B10, 05C38, 57N05

† Professor Voss passed away on September 7, 2003.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{T}\mathcal{E}\mathcal{X}$

4. Light subgraphs of order three	12
5. Subgraphs with restricted degrees of their vertices	15
6. Light subgraphs - maximum degree problems	16
7. Maximum degree of light families	19
8. The weight problems	20
9. Light subgraphs of graphs embedded on surfaces	23
10. Related topics	28
References	30

1. INTRODUCTION

The study of the structure of plane graphs (i.e. planar graphs embedded in the plane without edge crossings) has its origin in the time of L. Euler. It is connected with the lovely result (*Euler's polyhedron formula* discovered in 1750) stating that in a convex polyhedron with n vertices, e edges and f faces, $n - e + f = 2$. It was apparently discovered by Euler and first proved by Legendre (see, e.g. [BLW], [Mk]). The graph theory version of the formula is expressed in

Theorem 1.1 (Euler's Polyhedron Formula). *In a plane connected graph with n vertices, e edges and f faces,*

$$(1.1) \quad n - e + f = 2.$$

□

Euler's formula was not systematically exploited to any extent until the late nineteenth century. Only then a renaissance of interest in metric and combinatorial properties of solids started. A renewed interest in geometry and combinatorics of convex solids culminated in the landmark book [SR] by Steinitz and Rademacher published in 1934. This book presented an extraordinary result, known as *Steinitz' theorem*. Unfortunately, the theorem was couched in such an archaic language that it was not appreciated for many years. Only the reformulation of the theorem by Grünbaum in 1963 released a torrent of results. This resulted in a cross-fertilization of geometry with both graph theory and combinatorics, with benefits to all three areas. To be able to formulate this "*Fundamental Theorem on Polyhedral Graph Theory*" in modern terminology we need two definitions. The *graph of a polyhedron P* is the graph consisting of vertices and edges of P . A graph G is *polyhedral* if it is isomorphic to the graph of some convex polyhedron.

Theorem 1.2 (Steinitz' Theorem). *A graph is polyhedral if and only if it is planar and 3-connected.* □

This result is deeper than it might at first appear: proofs and excellent discussions may be found in ([Mk], [G1], [G3], [Zi]). What makes the theorem so remarkable is its implication that a general class of 3-dimensional structures is equivalent to a certain class of 2-dimensional ones - that is, studying convex polyhedra combinatorially does not require thinking of them in 3-dimensional space. It is sufficient to investigate their graphs. Hence any knowledge in 3-connected planar graphs indicates properties of convex polyhedra.

Probably the most important impulse to study the structure of plane graphs came from one of the most celebrated combinatorial problems - *The Four Colour Conjecture* (4CC), posed in 1852. This conjecture was proved in 1976 by Appel and Haken [AH] and the result is well known as

Theorem 1.3 (The Four Colour Theorem). *Vertices of every planar graph can be coloured with four colours in such a way that adjacent vertices are coloured with different colours.* \square

When trying to solve 4CC, Birkhoff in 1912 reviewed several ideas due to earlier writers, and welded them into a systematic method of investigation. The line of enquiry which he suggested led to the solution of the problem in 1976 (see Chapter 9 in [BLW]). This line is a common method for proofs of many theorems concerning properties of plane graph, see e.g. [AH], [Bo2], [Bo4], [Bo8], [Bo13], [BW2], [EHJ], [HM], [HJ2], [MS], [PT], [WL1].

If there are plane graphs which are counterexamples to a theorem, then there must be among them a graph with the smallest number of vertices; such a graph is said to be *irreducible* (or *minimal counterexample*) with respect to the theorem. The basic idea is to obtain more and more restrictive conditions which an irreducible graph must satisfy, in hope that eventually we shall have enough conditions either to construct the graph explicitly, or, alternatively, to prove that it cannot exist. These restrictive conditions are usually described in language of configurations.

A *configuration* H in a plane graph G is a connected portion of G (i.e. a connected subgraph H of G together with degrees of vertices of H in G). We call the set \mathcal{C} of configurations *unavoidable* in a given family \mathcal{G} of plane graphs if at least one member of \mathcal{C} is present in every graph from \mathcal{G} . As an example we note that the set of vertices of degree less than six is in this sense unavoidable in the family of all plane graph. We also define a configuration H to be *reducible* if it cannot be contained in any irreducible graph with respect to the conjecture.

Heawood in 1890 observed that if there was an irreducible plane graph with respect to the 4CC then it would belong to the set \mathcal{A} of plane triangulations of minimum degree 5. Hence to prove 4CC it is sufficient to find a finite unavoidable set \mathcal{U} of reducible configurations in \mathcal{A} . Appel and Haken [AH] were successful in finding such a set consisting of 1879 configurations (see also [HS], [WW]). In 1989 Appel and Haken [AH] announced that proofs of 4CC with only 1482, 1405, and 1256 configurations are possible. In an independent proof of 4CC Robertson, Sanders, Seymour, and Thomas [RSST] present an unavoidable set of 633 reducible configurations.

Already in 1904 Wernicke showed that every plane triangulation of minimum degree 5 contains either two adjacent vertices of degree 5 or a vertex of degree 5 adjacent to a vertex of degree 6, see [We]. In our modern language this reads as follows: "The set \mathcal{A} of all plane triangulations of minimum degree 5 has an unavoidable set of two configurations, namely, an edge with both end vertices of degree 5, and an edge with one end vertex of degree 5 and the second end vertex of degree 6."

In 1922 Franklin [Fr] extended Wernicke's result proving that each plane graph from the set \mathcal{A} contains a vertex of degree 5 adjacent either to two other vertices of degree 5, or to a vertex of degree 5 and a vertex of degree 6, or to two vertices of degree 6.

Attracted by 4CC and the result of Franklin, H. Lebesgue (the top personality of the *Mathematical Analysis* of the 20th century) realized that it would be very helpful to identify unavoidable sets of configurations for different families of plane graphs. In his 1940 paper [Le] he presented several such lists. His fundamental theorem (see Theorem 2.1 in Section 2 below) provides an unavoidable set of configurations for the family of all 3-connected plane graphs.

However, there is a limit to what can be achieved with Lebesgue's approach, and so stronger methods have been devised over the last three decades in order to solve various long-standing structural and colouring problems on plane graphs (for a survey of a portion of this area, see [Bo13]). As a result, many unavoidable sets of configurations for different families of plane graphs have been discovered. They are scattered in many papers. It would be an honourable achievement to collect and classify them.

We concentrate ourselves on unavoidable sets \mathcal{C} of graphs for given families \mathcal{G} of plane graphs that have the following property: Whenever a graph G from \mathcal{G} has a subgraph H from \mathcal{C} then it also contains a copy K of H such that every vertex of K has, in G , degree bounded by a constant $\varphi(\mathcal{C}, \mathcal{G})$ that depends only on the set \mathcal{C} and the family \mathcal{G} . We call the graphs from the family \mathcal{C} *light* in the family \mathcal{G} .

It is a well known fact easily deduced from Euler's Formula that each plane graph contains a vertex of degree at most five. Such a vertex can be interpreted as a path on one vertex. Thus such a path on one vertex is light in the class of all plane graphs. In 1955 A. Kotzig showed that each 3-connected plane graph contains an edge (e.i. a path with two vertices) of degree sum at most 13. We say that such paths with two vertices are light in the class of these graphs. The complete bipartite graphs $K_{2,s}$ for $s \geq 3$ show that the paths with two vertices are not light in the class of all plane graphs. In 1997 Fabrici and Jendrof proved that for each k the paths with k vertices are light in the family of all 3-connected plane graphs and no other plane graph H , different from a path is light in this family of graphs.

In this paper we give a survey of results on light subgraphs in several families of graphs embedded in the plane and the projective plane. Light subgraphs in graphs embedded in surfaces other than the plane or the projective plane are considered in another survey paper [JV10]. However, in Section 9 we briefly mention a recent progress together with the most important results concerning light subgraphs in graphs embedded in surfaces.

2. NOTATION AND PRELIMINARIES

All graphs considered throughout the paper have no loops or multiple edges. Multigraphs can have multiple edges and loops. An embedding of a planar graph (planar multigraph) into the plane \mathbb{M} is called a *plane graph* (a *plane multigraph*, respectively). If a planar (multi)graph G is embedded in \mathbb{M} , then the maximal connected regions of $\mathbb{M} - G$ are called the *faces* of G .

The *facial walk* of a face α of a connected plane multigraph G is the shortest closed walk traversing all edges incident with α . The *degree (or size)* of a face α is the length of its facial walk. The degree of a face α in G is denoted by $\deg_G(\alpha)$ or $\deg(\alpha)$ if G is known from the context. The degree of the vertex v of a connected plane multigraph G

is the number of incidences of edges with v , where loops are counted twice. Analogously, the notation $\deg_G(v)$ or $\deg(v)$ is used for the degree of a vertex v . Vertices and faces of degree i are called i -vertices and i -faces (or i -gons), respectively. The numbers of i -vertices and i -faces of a connected plane multigraph G are denoted $n_i(G)$ and $f_i(G)$, respectively, or n_i and f_i , if G is known. We use $\delta(G)$ to denote the minimum vertex degree of G .

We call an edge h an (a, b) -edge if the endvertices of h are an a -vertex and a b -vertex. By $e_{a,b}(G)$ or $e_{a,b}$ we denote the number of (a, b) -edges in a plane multigraph G .

An r -face α , $r \geq 3$, is said to be the (a_1, a_2, \dots, a_r) -face if vertices x_1, x_2, \dots, x_r , in order incident with α have degrees a_1, a_2, \dots, a_r . An (a_1, a_2, a_3) -face (an (a_1, a_2, a_3, a_4) -face, an $(a_1, a_2, a_3, a_4, a_5)$ -face) is called an (a_1, a_2, a_3) -triangle, (an (a_1, a_2, a_3, a_4) -quadrangle, or an $(a_1, a_2, a_3, a_4, a_5)$ -pentagon, respectively). Now we are able to state the classical theorem of Lebesgue [Le] already mentioned in Section 1

Theorem 2.1 (Lebesgue's Theorem). *Every 3-connected plane graph contains at least one of the following faces*

- (i) an (a, b, c) -triangle for

$a = 3$ and $3 \leq b \leq 6$ and $3 \leq c$, or	$a = 3$ and $b = 7$ and $7 \leq c \leq 41$, or
$a = 3$ and $b = 8$ and $8 \leq c \leq 23$, or	$a = 3$ and $b = 9$ and $9 \leq c \leq 17$, or
$a = 3$ and $b = 10$ and $10 \leq c \leq 14$, or	$a = 3$ and $b = 11$ and $11 \leq c \leq 13$, or
$a = 4$ and $b = 4$ and $4 \leq c$, or	$a = 4$ and $b = 5$ and $5 \leq c \leq 19$, or
$a = 4$ and $b = 6$ and $6 \leq c \leq 11$, or	$a = 4$ and $b = 7$ and $7 \leq c \leq 9$, or
$a = 5$ and $b = 5$ and $5 \leq c \leq 9$, or	$a = 5$ and $b = 6$ and $6 \leq c \leq 7$, or
- (ii) a $(3, b, c, d)$ -quadrangle for

$b = 3$ and $c = 3$ and $d \geq 3$, or	$b = 3$ and $c = 4$ and $4 \leq d \leq 11$, or
$b = 4$ and $c = 3$ and $4 \leq d \leq 11$, or	$b = 3$ and $c = 5$ and $5 \leq d \leq 7$, or
$b = 5$ and $c = 3$ and $5 \leq d \leq 7$, or	$b = 4$ and $c = 4$ and $4 \leq d \leq 5$, or
$b = 4$ and $c = 5$ and $d = 4$, or	
- (iii) a $(3, 3, 3, 3, d)$ -pentagon for $3 \leq d \leq 5$. □

Let $V(H)$ denote the set of vertices of a graph H . If H is a subgraph of a graph G , then the *weight* $w_G(H)$ of H in G is the sum of degrees in G of vertices of H .

$$w_G(H) = \sum_{v \in V(H)} \deg_G(v).$$

Moreover, the weight $w_G(e)$ of an (a, b) -edge e is

$$w_G(e) = a + b$$

If G is known from context, then we simply write $w(H) = w_G(H)$ and $w(e) = w_G(e)$.

A path and a cycle on k distinct vertices are described as a k -path and a k -cycle, respectively. A k -path is denoted by P_k . The *length* of a path or a cycle is the number of its edges. A k -path P_k with vertices v_1, v_2, \dots, v_k in order is also called an (a_1, a_2, \dots, a_k) -path, if $\deg(v_i) = a_i$ for all $i = 1, 2, \dots, k$.

Let $K_{1,3}$ be a subgraph of a graph G , we call it a $(d; a, b, c)$ -star if its central vertex has degree d and its three leaves have degrees a, b , and c in G .

For a connected plane multigraph G , let V, E , and F be the vertex set, the edge set, and the face set of G , respectively. Since

$$\sum_{\alpha \in F} \deg(\alpha) = \sum_{v \in V} \deg(v) = 2|E|,$$

from (1.1) we can easily derive

$$(2.1) \quad \sum_{\alpha \in F} (6 - \deg(\alpha)) + 2 \sum_{v \in V} (3 - \deg(v)) = 12$$

$$(2.2) \quad \sum_{v \in V} (6 - \deg(v)) + 2 \sum_{\alpha \in F} (3 - \deg(\alpha)) = 12$$

$$(2.3) \quad \sum_{v \in V} (4 - \deg(v)) + \sum_{\alpha \in F} (4 - \deg(\alpha)) = 8$$

Let $\mathcal{P}(\delta, \rho)$ be the family of all 3-connected plane graphs (i.e. polyhedral graphs, see Theorem 1.2) with minimum vertex degree at least δ and minimum face size at least ρ . $\mathcal{P}(\delta, \bar{\rho})$ stands for the family of all graphs from $\mathcal{P}(\delta, \rho)$ in which every face is a ρ -face. Analogously the family $\mathcal{P}(\bar{\delta}, \rho)$ is introduced. Note that the family $\mathcal{P}(3, \bar{3})$ is the family of all plane *triangulations* and $\mathcal{P}(3, \bar{4})$ is the family of all plane *quadrangulations*. By $\mathcal{P}(\delta, \rho; R)$ we mean a subfamily of $\mathcal{P}(\delta, \rho)$ the members of which fulfil the additional requirements R . It is an easy consequence of the equalities (2.1), (2.2) and (2.3) that $\mathcal{P}(\delta, \rho)$ is nonempty only when $(\delta, \rho) \in \{(3, 3), (3, 4), (4, 3), (3, 5), (5, 3)\}$.

Let $\mathcal{M}(\delta, \rho)$ be the family of all connected plane multigraphs with minimum vertex degree at least δ and minimum face size at least ρ . A plane multigraph G from the family $\mathcal{M}(3, 3)$ is called to be a *normal plane map*. If a plane multigraph from $\mathcal{M}(3, 3)$ has only 3-faces then it is called a plane *semitriangulation*. Note that semitriangulations can contain loops and multiple edges while triangulations do not have them.

Using (1.1) one can easily obtain

$$(2.4) \quad |E| \leq 3|V| - 6$$

for each normal plane map with $|V| \geq 3$.

For two graphs H and G we write $G \cong H$ if the graphs H and G are isomorphic. For a graph K we say that G contains a copy of K if G has a subgraph H such that $H \cong K$.

3. LIGHT EDGES

The theory of light subgraphs has its origin in two beautiful theorems of Kotzig [Ko1]. They state that every 3-connected plane graph contains an edge of weight (i.e. the sum of degrees of its endvertices) at most 13 in general, and at most 11 in the absence of 3-vertices, respectively. These bounds are best possible, as can be seen from the 3-connected plane

graphs obtained by placing 20 and 12 small pyramids on the faces of the icosahedron and dodecahedron, respectively, as well as for infinitely many other 3-connected plane graphs.

Kotzig's result was further developed in various directions. We shall discuss some of them in next sections. Here we only mention that in 1972 Erdős conjectured (see [G4]) that Kotzig's theorem is valid for all planar graphs with minimum vertex degree at least 3. This conjecture was proved (but never published) by Barnette (see [G4]) and independently by Borodin [Bo2]. The theorem of Kotzig was published in 1955 in Slovak. Therefore, its original proof [Ko1] is not readily accessible, but Grünbaum [G3] in 1975 gave a sketch of a proof in the English language. Other proofs can be deduced from [Bo2], [Bo3], [Bo5], [Bo10], [Je5]. Here we present a simple proof according to [Je4] that uses the Discharging method, a method used in the proof of the Four Color Theorem (see [AH], [RSST]). (Note that the idea of discharging is due to Heesch [H1]). This method is a common technique for proving results on planar graphs. We prove the theorem in a bit stronger form that implies the truth of Erdős' conjecture.

Theorem 3.1 (Kotzig's theorem). *Every normal plane map contains a $(3, a)$ -edge with $3 \leq a \leq 10$, or a $(4, b)$ -edge with $4 \leq b \leq 7$, or a $(5, c)$ -edge with $5 \leq c \leq 6$. The bounds 10, 7 and 6 are best possible.*

Proof. Let G be a counterexample on a set V of n vertices that has the maximum number of edges, say m , among all counterexamples on n vertices. Let f be the number of faces of G . For the purposes of the proof, edges of the desired type are called *light* edges.

By the choice of G , it must be a semitriangulation. Suppose G has a k -face α with $k \geq 4$. Because light edges are not present in G each edge has an endvertex of degree at least 6, so α is incident with two vertices x and y that are not consecutive on the boundary of α and both have degrees at least 6.

Inserting a diagonal xy into the face α we obtain a graph G^* having the same vertex set V as G but one edge more, a contradiction.

Because G is a semitriangulation, (2.2) may be rewritten

$$(3.1) \quad \sum_{v \in V} (\deg(v) - 6) = -12.$$

Consider an initial charge function $\varphi : V \rightarrow \mathbb{Q}$ such that $\varphi(v) = \deg(v) - 6$ for $v \in V$. Therefore (3.1) is equivalent to

$$\sum_{v \in V} \varphi(v) = -12.$$

We use the following rule in order to transform φ into a new charge function $\psi : V \rightarrow \mathbb{Q}$ by redistributing charges locally so that $\sum_{v \in V} \varphi(v) = \sum_{v \in V} \psi(v)$.

Rule. If $e = uv$ is an edge of G with $\deg(u) \geq 7$ and $\deg(v) \leq 5$. Then the vertex u sends along e to v the charge $\frac{6 - \deg(v)}{\deg(v)}$. Let $\psi(x)$ denote the resulting charge at a vertex x . Since charge sent to v is deducted from u , we have

$$\sum_{x \in V} \psi(x) = -12.$$

We are going to show that ψ is a nonnegative function, which will trivially be a contradiction. To this end consider several cases.

Case 1. Let v be a k -vertex for $3 \leq k \leq 6$. As G does not contain light edges, v receives a charge $\frac{6-k}{k}$ from each of its neighbours. Hence $\psi(v) = \varphi(v) + k\frac{6-k}{k} = k - 6 + 6 - k = 0$.

Case 2. Let u be a k -vertex for $k \geq 7$. Because G is a semitriangulation and does not contain light edges, at most half of the neighbors of u can have degree at most 5. Hence the vertex u sends a charge to at most $\lfloor \frac{k}{2} \rfloor$ vertices, all of degrees ≤ 5 .

2.1. $k = 7$. A transfer from u is possible only to 5-vertices, therefore

$$\psi(u) \geq \varphi(u) - \frac{1}{5} \left\lfloor \frac{7}{2} \right\rfloor = 7 - 6 - \frac{3}{5} > 0$$

2.2. $k \in \{8, 9, 10\}$. The vertex u sends charges only along $(k, 4)$ -edges or $(k, 5)$ -edges. Since charge $\frac{1}{2}$ would be sent along $(k, 4)$ -edges and only charge $\frac{1}{5}$ along $(k, 5)$ -edges we have

$$\psi(u) \geq \varphi(u) - \frac{1}{2} \left\lfloor \frac{k}{2} \right\rfloor = k - 6 - \frac{1}{2} \left\lfloor \frac{k}{2} \right\rfloor \geq 0.$$

2.3. If $k \geq 11$, then u sends charge $\frac{1}{5}$ along $(k, 5)$ -edges, charge $\frac{1}{2}$ along $(k, 4)$ -edges and charge 1 along $(k, 3)$ -edges. Since at most $\lfloor \frac{k}{2} \rfloor$ neighbours of u have degree at most 5, we obtain

$$\psi(u) \geq \varphi(u) - \left\lfloor \frac{k}{2} \right\rfloor = k - 6 - \left\lfloor \frac{k}{2} \right\rfloor \geq 0.$$

The bounds 10 and 6 are best possible as can be seen from the graphs mentioned in Section 2. An example showing that also 7 is best possible can be found in [Bo4].

This finishes the proof. \square

Now we turn our attention to the problem of estimating the number of light edges (i.e. the edges having weight at most 13) in families of plane graphs and multigraphs. Let $e_{i,j}(G) = e_{i,j}$ be the number of edges in a planar multigraph G that join i -vertices with j -vertices.

Theorem 3.1 of Kotzig states that $\sum_{i+j \leq 13} e_{i,j} > 0$ for every 3-connected planar graph. Grünbaum [G4] conjectured that for every 3-connected plane graph the following is true:

$$\begin{aligned} & 20e_{3,3} + 15e_{3,4} + 12e_{3,5} + 10e_{3,6} + \frac{20}{3}e_{3,7} + 5e_{3,8} + \frac{10}{3}e_{3,9} + 2e_{3,10} \\ & + 12e_{4,4} + 7e_{4,5} + 5e_{4,6} + 4e_{4,7} + \frac{8}{3}e_{4,8} + \frac{2}{3}e_{4,9} \\ & + 4e_{5,5} + 2e_{5,6} + \frac{1}{3}e_{5,7} + 12e_{6,6} \geq 120. \end{aligned}$$

Jucovič [Ju1] proved that each plane triangulation satisfies

$$\begin{aligned} &20e_{3,3} + 25e_{3,4} + 16e_{3,5} + 10e_{3,6} + \frac{20}{3}e_{3,7} + 5e_{3,8} + \frac{5}{2}e_{3,9} + 2e_{3,10} \\ &+ 20e_{4,4} + 11e_{4,5} + 5e_{4,6} + 6e_{4,7} + 5e_{4,8} + 3e_{4,9} \\ &+ 8e_{5,5} + 2e_{5,6} + 2e_{5,7} + 2e_{5,8} \geq 120. \end{aligned}$$

Later Jucovič [Ju2] proved that this inequality holds for all 3-connected planar graphs.

For the class of normal plane maps, which includes the class of 3-connected plane graphs, Borodin [Bo5] obtained the following result. (Recall that a normal plane map is a plane multigraph in which every vertex degree and every face size is at least 3.)

Theorem 3.2 [Bo5]. *For each normal plane map it holds that*

$$\begin{aligned} &40e_{3,3} + 25e_{3,4} + 16e_{3,5} + 10e_{3,6} + \frac{20}{3}e_{3,7} + 5e_{3,8} + \frac{5}{2}e_{3,9} + 2e_{3,10} \\ &+ \frac{50}{3}e_{4,4} + 11e_{4,5} + 5e_{4,6} + \frac{5}{3}e_{4,7} + \frac{16}{3}e_{5,5} + 2e_{5,6} \geq 120; \end{aligned}$$

Moreover, each coefficient of this inequality is best possible. \square

The sharpness of coefficients in Theorem 2 and in those below is understood in the sense that neither of coefficients may be decreased keeping all the other α_{ij} constant without violating the correspondent relation.

A final answer to the above mentioned Grünbaum's conjecture gives the following result by Fabrici and Jendrol' [FJ1]:

Theorem 3.3. [FJ1]. *For every 3-connected plane graph there is*

$$\begin{aligned} &20e_{3,3} + 25e_{3,4} + 16e_{3,5} + 10e_{3,6} + \frac{20}{3}e_{3,7} + 5e_{3,8} + \frac{5}{2}e_{3,9} + 2e_{3,10} \\ &+ \frac{50}{3}e_{4,4} + 11e_{4,5} + 5e_{4,6} + \frac{5}{3}e_{4,7} + \frac{16}{3}e_{5,5} + 2e_{5,6} \geq 120; \end{aligned}$$

Moreover, each coefficient is best possible. \square

The two inequalities for normal maps and 3-connected planar graphs, respectively, differ only in the coefficient of $e_{3,3}$. As was later proved by Borodin, Theorem 3.3 is valid for all normal plane maps with the exception of exactly one multigraph.

In the subclass of all normal plane maps with minimum degree at least 4 we have $e_{3,j} = 0$ for $3 \leq j \leq 10$. Borodin proved that in the resulting inequality all coefficients are best possible.

Theorem 3.4 [Bo5]. *For every normal plane map of minimum degree four, it holds that*

$$\frac{50}{3}e_{4,4} + 11e_{4,5} + 5e_{4,6} + \frac{5}{3}e_{4,7} + \frac{16}{3}e_{5,5} + 2e_{5,6} \geq 120$$

Moreover, each of these coefficients is best possible. \square

Already in 1904 in his work on the Four Color Problem Wernicke [We] has established the inequality $e_{5,5} + e_{5,6} > 0$ for every graph $G \in \mathcal{P}(5, 3)$. Further contributions are due to Grünbaum [G2], [G4], Fisk (see [GS]), Grünbaum and Shephard [GS], and Borodin [Bo6]. Borodin and Sanders [BS] found the best possible light edge inequality for plane graphs of minimum degree five.

Theorem 3.5 [BS]. *For every normal plane map of minimum vertex degree at least 5 and minimum face size 3 it holds that*

$$\frac{14}{3}e_{5,5} + 2e_{5,6} \geq 120.$$

Moreover, the coefficients $\frac{14}{3}$ and 2 are best possible. \square

If in Theorem 3.5 we use $e_{4,j} = 0$ for $4 \leq j \leq 7$ then the resulting inequality differs from that of Theorem 3.4 only in the coefficient of $e_{5,5}$.

We finish this section with few very recent results.

A planar graph which can be embedded in the plane in such a way that every vertex lies on the boundary of the same region is called an *outerplanar* graph. Hackmann and Kemnitz [HK] recently proved

Theorem 3.6 [HK]. *Every outerplanar graph with minimum degree at least 2 contains an $(2, b)$ -edge for $2 \leq b \leq 3$ or a $(2, 4, 2)$ -path. \square*

Theorem 3.7 [JM]. *Every planar graph of minimum degree five contains a $(5; a, b, c)$ -star for $a = 5$ and $5 \leq b \leq 6$ and $5 \leq c \leq 7$, or $a = b = c = 6$.*

Moreover the bounds 6 and 7 are best possible. \square

The next theorem is a strengthening of Theorems 3.1 and 3.7.

Theorem 3.8 [FHJ]. *Every planar graph G of minimum degree at least 3 contains*

- (i) a $(3, b)$ -edge for $3 \leq b \leq 10$, or
- (ii) an $(a, 4, b)$ -path for
 - $a = 4$ and $4 \leq b \leq 10$, or
 - $a = 5$ and $5 \leq b \leq 9$, or
 - $6 \leq a \leq 7$ and $6 \leq b \leq 8$, or
- (iii) a $(5; a, b, c)$ -star for
 - $4 \leq a \leq 5$ and $5 \leq b \leq 6$ and
 - $5 \leq c \leq 7$, or $a = b = c = 6$.

Moreover, for every $S \in \{(3, 10)\text{-edge}, (4, 4, 9)\text{-path}, (5, 4, 8)\text{-path}, (6, 4, 8)\text{-path}, (7, 4, 7)\text{-path}, (5; 5, 6, 7)\text{-star}, (5; 6, 6, 6)\text{-star}\}$ there is a 3-connected plane graph H containing S and no other configuration from the above list. \square

Theorem 3.8 has an application in a problem of colouring of vertices of the square of planar graph posed by Wegner [Wg] in 1977 (see also Jensen and Toft [JT], p.51). Recall that the square G^2 of a graph G is a graph with the same vertex set $V(G)$. Two vertices x and y are adjacent in G^2 if and only if their distance in G is at most 2.

Corollary 3.8.1 [FHJ]. *Let G be a planar graph of maximum degree $\Delta \geq 11$. Then for the chromatic number $\chi(G^2)$ of the square of G there holds:*

$$\chi(G^2) \leq 2\Delta + 19.$$

Proof. Suppose there is a counterexample. Let H be one with the minimum number of vertices. Evidently the minimum degree of H is at least 3. By Theorem 3.8 the graph H contains an (a, b) -edge $e = xy$ with $\deg_H(x) = a$, $\deg_H(y) = b$ and $\deg_{H^2}(x) \leq 2\Delta + 18$ with values $a = 3$ and $b \leq 10$, or $a = 4$ and $b \leq 7$, or $a = 5$ and $b \leq 6$, and H^2 being the square of H . If the edge e is contracted into a vertex z and all multiple edges are simplified, the resulting graph H^* has the same maximum degree but it is not a counterexample. The square of H^* has a colouring with $2\Delta + 19$ colours. This colouring induces a colouring with $2\Delta + 19$ colours of the graph H in which all vertices except of x and y are coloured. Then the colour of vertex z is assigned to the vertex y . As the vertex x has, in H^2 , at most $2\Delta + 18$ neighbours it can be coloured with one of the available $2\Delta + 19$ colours, a contradiction. \square

Note that the bound of Corollary 3.8.1 is better than $2\Delta + 25$, the bound obtained recently by van den Heuvel and McGuinness [HM].

The requirement on minimum degree at least 3 in the above theorems cannot be relaxed as one can see from the graphs $K_{1,r}$ and $K_{2,r}$ for $r \geq 3$.

Theorem 3.9 [ABG]. *Every connected planar graph of order at least 2 contains either*

- (i) *two vertices having degree sum ≤ 4 , or*
- (ii) *two 3-vertices at distance two, or*
- (iii) *an edge e of weight at most 11 incident with two 3-faces, or*
- (iv) *an edge g of weight at most 9 incident with a 3-face, or*
- (v) *an edge h of weight at most 7.*

Moreover the bounds 11, 9, and 7 are tight. \square

For other results in this direction, see Borodin [Bo3], [Bo5], [Bo6], [Bo13], Borodin and Sanders [BS], Fabrici, Harant, and Jendroľ [FHJ], Jucovič [Ju2], Sanders [Sa], and Zaks [Za1], [Za2]. Note that a result weaker than the theorem of Kotzig was known already to Lebesgue. In 1940 he proved that every $G \in \mathcal{P}(3, 3)$ contains an edge e with $w(e) \leq 14$, see [Le] and Theorem 2.1.

4. LIGHT SUBGRAPHS OF ORDER THREE

Trying to solve the Four Color Map Problem Franklin [Fr] in 1922 proved that every 3-connected plane graph G of minimum degree at least 5, contains a 3-path P_3 with weight 17, the bound being best possible. Only recently, in 1993, Ando, Iwasaki, and Kaneko obtained the analogous result for all 3-connected plane graphs. Namely, they proved

Theorem 4.1 ([AIK]). *Every $G \in \mathcal{P}(3,3)$ contains a 3-path P_3 with weight at most 21. Moreover the bound 21 is sharp.*

The following result has been proved by Jendrof [Je2] which strengthens Theorem 3.1 of Kotzig [Ko1].

Theorem 4.2 ([Je2]). *Every $G \in \mathcal{P}(3,3)$ contains an (a,b,c) -path where*

- (i) $3 \leq a \leq 10$ and $b = 3$ and $3 \leq c \leq 10$, or
- (ii) $4 \leq a \leq 7$ and $b = 4$ and $4 \leq c \leq 7$, or
- (iii) $5 \leq a \leq 6$ and $b = 5$ and $5 \leq c \leq 6$, or
- (iv) $a = 3$ and $3 \leq b \leq 4$ and $4 \leq c \leq 15$, or
- (v) $a = 3$ and $5 \leq b \leq 6$ and $4 \leq c \leq 11$, or
- (vi) $a = 3$ and $7 \leq b \leq 8$ and $4 \leq c \leq 5$, or
- (vii) $a = 3$ and $3 \leq b \leq 10$ and $c = 3$, or
- (viii) $a = 4$ and $b = 4$ and $4 \leq c \leq 11$, or
- (ix) $a = 4$ and $b = 5$ and $5 \leq c \leq 7$, or
- (x) $a = 4$ and $6 \leq b \leq 7$ and $4 \leq c \leq 5$.

Moreover, in each of the cases (iii), (iv), (vi), and (vii) the upper bounds on parameters a, b, c can be obtained simultaneously. Furthermore, there is a graph in $\mathcal{P}(3,3)$ having only $(4,7,4)$ -paths and $(7,4,7)$ -paths from the above list. \square

The requirement of 3-connectedness in the above theorems is fundamental because of the following theorem. We present a proof of this theorem with a proof technique that is typical in this area.

Theorem 4.3 [Je6]. *For every connected plane graph H of order at least 3 and every integer $m \geq 3$ there exists a 2-connected plane graph G such that each copy of H in G contains a vertex A with*

$$\deg_G(A) \geq m$$

Proof. Augment H to a triangulation T with vertex set $V(H)$. Let $[uvw]$ be an outer 3-face of T . For $m \geq 2$, let D_m be a plane graph obtained from the double $2m$ -pyramid with poles z_1 and z_2 by deleting every second edge of the equatorial cycle C_{2m} . If we insert into a 3-face $[z_1xy]$ of D_m the triangulation T so that the vertex u coincides with r_1 , the vertices v and w with x and y , respectively, we obtain the required graph G . If H has at least three vertices then each copy of H in G contains at least one of the vertices z_1 or z_2 , which have degree at least $2m$. \square

The graph G from the proof of Theorem 4.3 contains a 3-face incident with two 3-vertices. Borodin [Bo11] observed that if such faces are excluded the requirement of 3-connectivity can be omitted. More precisely he has proved.

Theorem 4.4 [Bo11]. *Each normal plane map G having no 3-faces incident with two 3-vertices has the following two properties:*

- (i) G has either a 3-path P_3 with $w(P_3) \leq 18$ or a vertex of degree ≤ 15 adjacent to two 3-vertices
- (ii) G has either a 3 path P'_3 with $w(P'_3) \leq 17$ or an edge e with $w(e) \leq 7$. \square

Already in 1940 Lebesgue [Le] proved that each 3-connected plane graph G of minimum degree 5 contains a 3-cycle C_3 bounding a 3-face α with $w(C_3) \leq 19$. Kotzig [Ko2] improved this result to $w(C_3) \leq 18$ and conjectured in [Ko3] that 17 is the best upper bound. Borodin confirms this in

Theorem 4.5 [Bo1]. *Every $G \in \mathcal{P}(5, 3)$ contains a 3-face α with $w_G(\alpha) \leq 17$. This bound is best possible.* \square

This theorem has a beautiful corollary. It confirms the conjecture of Grünbaum from 1975, see [G3], that every 5-connected plane graph is cyclically 11-connected. Let us sketch a proof of this statement. First recall that a graph is *cyclically k -connected* for some $k \geq 1$ if there is no set S of less than k -edges with property that $G - S$ has two components each containing a cycle. Let G be a graph from $\mathcal{P}(5, 3)$. Consider a face α of weight at most 17 of G . Choose S to be the set of edges incident with vertices of α but not on the boundary of α . Clearly $|S| \leq 11$ and the graph $G - S$ has two components each containing a cycle.

Note that already earlier, in 1972, Plummer [P] proved that the cyclic connectivity of these graphs is at most 13.

Many papers have studied the structural properties of different classes of plane triangulations, see e.g. [Ko2], [Ko3], Borodin [Bo6], [Bo9], [Bo12], [BB], Jendrol' [Je5], Sanders [Sa], Borodin and Sanders [BS]. Recently Jendrol' [Je5] proved the following theorem which includes earlier results by Lebesgue [Le], Kotzig [Ko2], [Ko3] and Borodin [Bo9].

Theorem 4.6 [Je5]. *Each plane triangulation of order at least five contains an (a, b, c) -triangle, where*

- (i) $a = 3$ and $b = 4$ and $4 \leq c \leq 35$, or
- (ii) $a = 3$ and $b = 5$ and $5 \leq c \leq 21$, or
- (iii) $a = 3$ and $b = 6$ and $6 \leq c \leq 20$, or
- (iv) $a = 3$ and $b = 7$ and $7 \leq c \leq 16$, or
- (v) $a = 3$ and $b = 8$ and $8 \leq c \leq 14$, or
- (vi) $a = 3$ and $b = 9$ and $9 \leq c \leq 14$, or
- (vii) $a = 3$ and $b = 10$ and $10 \leq c \leq 13$, or
- (viii) $a = 4$ and $b = 4$ and $c \geq 4$, or
- (ix) $a = 4$ and $b = 5$ and $5 \leq c \leq 13$, or
- (x) $a = 4$ and $b = 6$ and $6 \leq c \leq 17$, or
- (xi) $a = 4$ and $b = 7$ and $7 \leq c \leq 8$, or
- (xii) $a = 5$ and $b = 5$ and $5 \leq c \leq 7$, or
- (xiii) $a = 5$ and $b = 6$ and $c = 6$.

Moreover, if $c(k)$ denotes the upper bound on c in the above case (k), then $c(\text{i}) \geq 30, c(\text{ii}) \geq 18, c(\text{iii}) = 20, c(\text{iv}) \geq 7, c(\text{v}) = 14, c(\text{vi}) \geq 11, c(\text{vii}) \geq 12, c(\text{viii}) = \infty, c(\text{ix}) \geq 10, c(\text{x}) \geq 10, c(\text{xi}) \geq 7$ and $c(\text{xii}) = 7$. \square

Note that Theorem 4.6 considers all cases since plane triangulations have no multiple edges and hence every plane triangulation with at least five vertices has no $(3, 3)$ -edge.

Theorem 4.6 points out that if a plane triangulation has no $(4, 4)$ -edge, then it contains a light triangle. Borodin [Bo12] has gone further. He proved

Theorem 4.7 [Bo12]. *If in a plane triangulation T there is no path consisting of k vertices of degree 4 for some $k \geq 1$, then*

(i) *T contains a 3-face α with weight*

$$w(\alpha) \leq \max\{37, 10k + 17\} \text{ and}$$

(ii) *either there is a $(3, 4)$ -edge or a 3-face β with weight*

$$w(\beta) \leq \max\{29, 5k + 8\},$$

where all bounds are best possible. □

In a connection with results mentioned in this section, the following problem seems to be interesting.

Problem 4.1. *Find the best versions of Theorem 4.2. and Theorem 4.6*

In each of the cases (iii), (iv), (vi) and (vii) of Theorem 4.2 the upper bounds on parameters are tight and can be obtained independently from the others. Furthermore, there is a graph $G \in \mathcal{P}(3, 3)$ having only $(4, 7, 4)$ -paths and $(7, 4, 7)$ -paths from the list of the Theorem.

Similarly we are interested in the best version of Theorem 4.6. As we have mentioned, in the cases (iii), (v), (viii), (xii) and (xiii) the assertions of Theorem 4.6 are best possible. For the other cases we believe the following Conjecture is true:

Conjecture 4.1 [Je4]. *If $c(k)$ denotes the upper bound on c in the case (k) of Theorem 4.6, then $c(\text{i})=30$, $c(\text{ii})=18$, $c(\text{iv})=14$, $c(\text{vi})=2$, $c(\text{vii})=12$, $c(\text{ix})=10$ and $c(\text{xi})=7$. □*

For a plane graph G from $\mathcal{P}(3, 3)$, let $f_{i,j,k}$ be the number of 3-faces that are incident with an i -vertex, a j -vertex, and a k -vertex. As proved by Lebesgue [Le], $f_{5,5,5} + \frac{2}{3}f_{5,5,6} + \frac{3}{7}f_{5,5,7} + \frac{1}{4}f_{5,5,8} + \frac{1}{9}f_{5,5,9} + \frac{1}{3}f_{5,6,6} + \frac{2}{21}f_{5,6,7} \geq 120$ holds for every graph $G \in \mathcal{P}(5, 3)$. This result of Lebesgue and Theorem 4.5 are strengthened in the next Theorem proved by Borodin [Bo7] except fact that the optimality of the coefficient 5 of $f_{5,5,7}$. The best possibility of it was proved by Borodin and Sanders in [BS].

Theorem 4.8. *If G is a graph from $\mathcal{P}(5, 3)$, then*

$$18f_{5,5,5} + 9f_{5,5,6} + 5f_{5,5,7} + 4f_{5,6,6} \geq 144.$$

Moreover, all of these coefficients are best possible. □

5. SUBGRAPHS WITH RESTRICTED DEGREES

In this section we prove two basic results that are typical for this topic and have served as a starting point for a theory of light subgraphs which we shall describe in further sections. Because of Theorem 4.3 we only deal with the family of 3-connected plane graphs. The first result is due to Fabrici and Jendrol'.

Theorem 5.1 [FJ2]. *If G is a 3-connected plane graph having a k -path where k is a positive integer, then G contains a k -path P_k such that all vertices of P_k have degree at most $5k$ in G . Moreover, the bound $5k$ is tight.*

Proof. (a) *The upper bound.* Suppose the theorem is not true, and let G be a counterexample on n vertices that has the most edges among all counterexamples on n vertices. Let us call a vertex *major* (*minor*) if its degree is $> 5k$ ($\leq 5k$, respectively). The crucial point of the proof is in the following

Claim (*) *Every major vertex is incident only with triangular faces.*

Otherwise, there is a major vertex u incident with a face α of size at least 4. A diagonal uv can be inserted into α that joins u with a vertex v of α not adjacent with u . The graph G^* so obtained is again a counterexample, because the edge AB cannot appear on any k -path consisting of minor vertices only. But G^* has one edge more than G , which contradicts the maximality of G . This completes the proof of Claim (*).

Let M be a subgraph of G induced by major vertices of G . Since M is also a plane graph, it contains a vertex x such that

$$(5.1) \quad \deg_M(x) \leq 5.$$

On the other hand, because x is major, we have

$$\deg_G(x) \geq 5k + 1.$$

Due to Claim (*) there is a subgraph of G on the vertex x and its neighbours that is a wheel with $\deg_G(x)$ spokes. The ends of spokes different from x form a cycle with at least $5k + 1$ vertices. Among them there are at most five major vertices, by (5.1). By the pigeonhole principle, there is a path on the wheel consisting of k minor vertices, a contradiction.

(b) *The sharpness of the bound.* Now we shall construct a 3-connected plane graph G in which every k -path has a vertex u of degree at least $5k$. The construction begins with the dodecahedron. Into each of its 5-faces we insert a new vertex x and join it to all vertices incident with this face. The result is a graph all faces of which are triangles $[xyz]$ with $\deg(x) = 5$ and $\deg(y) = \deg(z) = 6$. Into every triangle of this graph we insert a subdivided 3-star consisting of a central vertex v and three paths p_x, p_y, p_z from v to x, y, z , respectively, with p_x of length $\lceil \frac{k+2}{2} \rceil$ and p_y and p_z of length $\lfloor \frac{k-2}{2} \rfloor$. Now make x adjacent to all vertices of p_x and p_y , and similarly make y adjacent to all of p_y and p_z , and z adjacent to all of p_z and p_x . Observe that in the resulting graph G all the vertices x have degree $5k$, the vertices y and z have degrees at least $6k - 6$ and every other vertex has degree at most 6. It is easy to see that each k -path contains at least one vertex of type x, y or z . \square

A natural question arises whether every 3-connected graph G having a copy of a connected graph H different from a path must also contain a copy of H such that its vertices have bounded degrees in G . The answer is surprisingly negative. Fabrici and Jendrof proved

Theorem 5.2 [FJ2]. *If H is a connected plane graph other than a path and m is an integer greater than $|V(H)|$. Then there is a 3-connected planar graph T such that each copy of H in T has a vertex y with*

$$\deg_T(y) \geq m.$$

Proof. Augment H to a triangulation T_o with vertex set $V(H)$. Into each triangle $[uvw]$ of T_o insert a wheel with a central vertex z and with m spokes zx_i for $1 \leq i \leq m$. Join the vertex u to x_i for $1 \leq i \leq \lfloor \frac{m}{3} \rfloor$, the vertex v to x_j for $\lfloor \frac{m}{3} \rfloor \leq j \leq \lfloor \frac{2m}{3} \rfloor$, and the vertex w to x_1 and to x_t for $\lfloor \frac{2m}{3} \rfloor \leq t \leq m$. The result is the desired triangulation T . One can easily check that each vertex of T that lies in T_o has degree at least m in T , as does the center of each inserted wheel. The vertices of degrees less than k induce m -cycles in T . Therefore, as H is not a path, each copy of H in T contains at least one vertex of degree $\geq m$. \square

Recently Madaras improved Theorem 5.1 by showing the following result.

Theorem 5.3 [Ma1]. *If G is a 3-connected plane graph containing a vertex of degree at least k , where k is a positive integer, then G contains a k -path P_k such that $k - 1$ vertices of P_k have degree at most $\frac{5}{2}k$, and the remaining vertex has degree at most $5k$ in G . \square*

6. MAXIMUM DEGREE PROBLEMS

The problems mentioned in the previous sections suggest the formulation of more general problems.

Problem 6.1. *Let \mathcal{H} be a family of graphs and let H be a connected graph that is a proper subgraph of at least one member of \mathcal{H} . Let $\varphi(H, \mathcal{H})$ be the smallest integer k with the property that every graph G in \mathcal{H} , that contains H contains a copy of H whose vertices all have degree at most k in G . Determine the value of $\varphi(H, \mathcal{H})$ for given H and \mathcal{H} .*

If such a $\varphi(H, \mathcal{H})$ does not exist then we write $\varphi(H, \mathcal{H}) = +\infty$. If $\varphi(H, \mathcal{H}) < +\infty$, then we call the graph H *light* in the family \mathcal{H} .

Here we consider the family $\mathcal{H} = \mathcal{P}(\delta, \rho)$ of all 3-connected plane graphs (i.e. the family of all polyhedral graphs [G1]) with minimum vertex degree at least δ and minimum face size at least ρ , where δ and ρ are at least 3. In the sequel, let

$$\varphi(\delta, \rho; H) = \varphi(H, \mathcal{P}(\delta, \rho))$$

and

$$\varphi(\delta, \bar{\rho}, H) = \varphi(H, \mathcal{P}(\delta, \bar{\rho})).$$

The result for P_1 derived from Euler's formula can be rewritten as $\varphi(3, 3; P_1) = 5$. Kotzig [Ko1] (see also Section 3) proved that each graph $G \in \mathcal{P}(3, 3)$ contains an edge e with $w(e) \leq 13$. This implies $\varphi(3, 3; P_2) = 10$. Theorem 4.1 provides $\varphi(3, 3; P_3) = 15$. Fabrici and Jendrol' (see Section 5) generalized the results for P_1, P_2 and P_3 to arbitrary P_k . Their Theorem 5.1 states that the path P_k is light in the family $\mathcal{P}(3, 3)$ for every k . Next theorem summarizes results concerning the constant $\varphi(P_k, \mathcal{G})$ for several families of 3-connected plane graphs \mathcal{G} . (Brackets indicate the papers, where the results are proved.)

Theorem 6.1.

- (i) $\varphi(3, 3; P_k) = 5k$ for all $k \geq 1$. [FJ1]
 - (ii) $\varphi(4, 3; P_k) = 5k - 7$ for all $k \geq 8$. [FHJW]
 - (iii) $5 \lfloor \frac{k}{2} \rfloor \leq \varphi(3, 4; P_k) \leq \frac{5}{2}k$ for all $k \geq 2$. [HJT]
 - (iv) $5k - 325 \leq \varphi(5, 3; P_k) \leq 5k - 7$ for all $k \geq 68$. [Fa1]
 - (v) $\frac{5}{3}k - 80 \leq \varphi(3, 5; P_k) \leq \frac{5}{3}k$ for all $k \geq 935$. [JO]
 - (vi) $2k + 2 \leq \varphi(P_k, \mathcal{F}) \leq 2k + 3$ for all $k \geq 2$. [HW]
- (Here \mathcal{F} is the family of 4-connected planar graphs) □

Note that the precise values of $\varphi(4, 3; P_k)$ are known also for all $k \leq 7$, see [FHJW]. We can also show that the lower bound in (iv) cannot be smaller than $5k - 125$ if $k \geq 714$ and that the upper bound in (v) is valid for all $k \geq 2$.

Theorem 5.2 asserts that no connected planar graph distinct from a k -path P_k for every k is light in the family $\mathcal{P}(3, 3)$. This motivates the following

Problem 6.2. For a given infinite family \mathcal{G} of plane graphs determine all connected planar graphs that are light in \mathcal{G} . □

The next theorem provides results of this type.

Theorem 6.2. If $\mathcal{G} \in \{\mathcal{P}(3, 3), \mathcal{P}(3, 4), \mathcal{P}(4, 3), \mathcal{F}\}$, where \mathcal{F} is the family of 4-connected planar graphs, then a graph H is light in the family \mathcal{G} if and only if H is k -path P_k for some $k \geq 1$. □

The proof of Theorem 6.2 for the family $\mathcal{P}(3, 4)$ is by Harant, Jendrol' and Tkáč [HJT], for $\mathcal{P}(4, 3)$ by Fabrici, Hexel, Jendrol' and Walther [FHJW], and for the family \mathcal{F} by Mohar [Mo1].

In the graph families $\mathcal{P}(3, 3), \mathcal{P}(3, 4)$, and $\mathcal{P}(4, 3)$ only the paths P_k are light. The situation changes significantly in the families $\mathcal{P}(5, 3)$ and $\mathcal{P}(3, 5)$. From Theorem 6.1 it follows that each path $P_k, k \geq 1$, is light in both families $\mathcal{P}(5, 3)$ and $\mathcal{P}(3, 5)$.

Next theorem excludes some families of graphs to be light

Theorem 6.3 [Fa1], [JMST], [JO].

- (i) No plane connected graph H with maximum degree at least 5 or with a block having at least 11 vertices is light in $\mathcal{P}(5, \bar{3})$ (and, hence, in $\mathcal{P}(5, 3)$).
- (ii) No plane connected graph H with maximum degree at least 4 or with a block having at least 19 vertices is light in $\mathcal{P}(3, 5)$. □

From the classical result of Lebesgue [Le] (see Theorem 2.1) it follows that the 3-cycle C_3 is light in $\mathcal{P}(5, 3)$ and the 5-cycle C_5 is light in $\mathcal{P}(3, 5)$. Recently Jendrol' and Madaras in

[JM] proved that the star $K_{1,r}$ for $r \geq 3$ is light in $\mathcal{P}(5, 3)$ if and only if $r \in \{3, 4\}$. However, the problem of determining all graphs that are light in $\mathcal{P}(5, 3)$ and $\mathcal{P}(3, 5)$ remains open. Jendrof et al. [JMST] proved the following theorem

Theorem 6.4 [JMST]. *The r -cycle C_r is light in the family $\mathcal{P}(5, \bar{3})$ of all plane triangulations with minimum degree 5 if and only if $3 \leq r \leq 10$. Moreover*

$$\begin{aligned} \varphi(C_3) &= 7, & \varphi(C_4) &= 10, & \varphi(C_5) &= 10, \\ 10 \leq \varphi(C_6) \leq 11, & 15 \leq \varphi(C_7) \leq 17, & 15 \leq \varphi(C_8) \leq 29, \\ 19 \leq \varphi(C_9) \leq 41, & 20 \leq \varphi(C_{10}) \leq 415, \end{aligned}$$

where $\varphi(C_k) = \varphi(5, \bar{3}; C_k)$. □

In [He] there is a question for which $k \geq 11$ the cycle C_k is light in the family of all 5-connected plane triangulation. The answer is rather surprising

Theorem 6.5 [HS]. *The k -cycle C_k is light in the family of all 5-connected plane triangulation for every k at least 3.* □

Hexel and Soták conjecture that for every $k \geq 3$ the k -cycle C_k is light in the family of all 5-connected planar graphs. More results concerning light graphs in subfamilies of plane graphs with high connectivity can be found in [Mo1] and [HW].

An analogue of Theorem 6.4 for the family $\mathcal{P}(3, \bar{5})$ is

Theorem 6.6 [JO]. *The r -cycle C_r is light in the family $\mathcal{P}(3, \bar{5})$ if and only if $r \in \{5, 8, 11, 14\}$. Moreover $\varphi(3, \bar{5}; C_5) = 5$, $6 \leq \varphi(3, \bar{5}; C_8) \leq 7$, $10 \leq \varphi(3, \bar{5}; C_{11}) \leq 11$ and $10 \leq \varphi(3, \bar{5}; C_{14}) \leq 17$.* □

By Theorem 6.6 the only cycles that are candidates for being light in $\mathcal{P}(3, 5)$ are C_5, C_8, C_{11} and C_{14} . As we mentioned above, C_5 is light in this family. In [JO] the cycles C_8 and C_{11} are proved to be not light there. It is an open question whether C_{14} is light in $\mathcal{P}(3, 5)$.

The 3-cycle is light in $\mathcal{P}(5, 3)$ and not light in $\mathcal{P}(4, 3)$. The question arises for which subclasses other than $\mathcal{P}(5, 3)$ the 3-cycle is light? Let $\mathcal{P}(4, 3, \mathcal{E}_t)$ denote the family of all graphs in $\mathcal{P}(4, 3)$ having no path consisting of t vertices all having degree 4. Borodin [Bo12] showed that the triangle C_3 is light in $\mathcal{P}(4, \bar{3}, \mathcal{E}_t)$ for all $t \geq 1$. Mohar, Škrekovski, and Voss [MŠV] showed that the cycle C_4 is light in $\mathcal{P}(4, 3, \mathcal{E}_t)$ for $t \in \{2, 3, 4\}$; so C_4 is light in $\mathcal{P}(4, 3, \mathcal{E}_t)$ for $1 \leq t \leq 4$. Further, C_4 is not light in $\mathcal{P}(4, 3, \mathcal{E}_t)$ for all $t \geq 23$. For $5 \leq t \leq 22$ this question is open. For $r \geq 5$ and $t \geq 3$ the cycle C_r is not light in $\mathcal{P}(4, 3, \mathcal{E}_t)$. For $t = 2$ Mohar, Škrekovski, and Voss showed

Theorem 6.7 [MŠV]. *The r -cycle C_r is light in the family $\mathcal{P}(4, 3, \mathcal{E}_2)$ of all 3-connected plane graphs of minimum degree at least 4 and edge-weight at least 9 if and only if $r \in \{3, 4, 5, 6\}$. Moreover,*

$$\begin{aligned} \varphi(4, 3, \mathcal{E}_2; C_3) &= 12, & \varphi(4, 3, \mathcal{E}_2; C_4) &\leq 22, \\ \varphi(4, 3, \mathcal{E}_2; C_5) &\leq 107, & \varphi(4, 3, \mathcal{E}_2; C_6) &\leq 107. \end{aligned}$$

□

From a theorem of Borodin [Bo1] follows: $\varphi(5, 3; C_3) = 7$. Soták (oral communication) proved $\varphi(5, 3; C_4) = 11$ and $\varphi(5, 3; C_5) = 10$. The proof that $\varphi(5, 3, C_6) \leq 107$ is by Mohar et al. [MŠV]. The lightness of the 7-cycle in $\mathcal{P}(5, 3)$ is proved by Madaras et al. [MaŠV]. Together with Theorem 6.7, these results imply

Theorem 6.8. *The r -cycle C_r is light in $\mathcal{P}(5, 3)$ if $r \in \{3, 4, 5, 6, 7\}$, and is not light in $\mathcal{P}(5, 3)$ if $r \geq 11$. Moreover,*

$$\varphi(5, 3; C_3) = 7, \varphi(5, 3; C_4) = 11, \varphi(5, 3; C_5) = 10, \varphi(5, 3; C_6) \leq 107, \varphi(5, 3; C_7) \leq 359. \quad \square$$

It is an open question whether in the class $\mathcal{P}(5, 3)$ the cycles C_8, C_9, C_{10} are light or not.

Jendroľ and Madaras [JM] showed for $r \geq 3$ that the star $K_{1,r}$ is light in $\mathcal{P}(5, 3)$ if and only if $r \in \{3, 4\}$. Mohar, Škrekovski, and Voss [MŠV] proved that this is also true in the class $\mathcal{P}(4, 3, \mathcal{E}_t)$. For $t \geq 3$ the only star that is light in $\mathcal{P}(4, 3, \mathcal{E}_t)$ is $K_{1,3}$, see [MŠV].

7. MAXIMUM DEGREE OF LIGHT FAMILIES

In section 6 we have defined a light subgraph H in a family \mathcal{H} of graphs. Here we introduce the concept of a light class \mathcal{L} of graphs in a family \mathcal{H} of graphs.

Problem 7.1. *Let \mathcal{H} be a family of graphs, and let \mathcal{L} be a finite family of connected graphs having the property that every member of \mathcal{L} is a proper subgraph of at least one member of \mathcal{H} . Let $\varphi(\mathcal{L}, \mathcal{H})$ be the smallest integer t with the property that every graph G in \mathcal{H} that has a subgraph belonging to \mathcal{L} , has such a subgraph whose vertices all have degree at most t in G . Determine the value $\varphi(\mathcal{L}, \mathcal{H})$ for a given pair of families \mathcal{L} and \mathcal{H} .*

If such a $\varphi(\mathcal{L}, \mathcal{H})$ does not exist, then we write $\varphi(\mathcal{L}, \mathcal{H}) = +\infty$. If $\varphi(\mathcal{L}, \mathcal{H}) < +\infty$, then we call the family \mathcal{L} *light* in the family \mathcal{H} . When \mathcal{L} is the family \mathcal{T}_k of all trees on k vertices, and $\mathcal{H} = \mathcal{P}(\delta, \rho)$, we write $\tau(k, \delta, \rho)$ for $\varphi(\mathcal{T}_k, \mathcal{P}(\delta, \rho))$.

Obviously, $\mathcal{T}_1 = \{P_1\}$, $\mathcal{T}_2 = \{P_2\}$, $\mathcal{T}_3 = \{P_3\}$, and $\{P_k\} \subset \mathcal{T}_k$ for all $k \geq 4$. Hence $\tau(k, \delta, \rho) = \varphi(\delta, \rho; P_k)$ for $1 \leq k \leq 3$, and $\tau(k, \delta, \rho) \leq \varphi(\delta, \rho; P_k)$. For $\mathcal{P}(3, 3)$ Fabrici and Jendroľ proved:

Theorem 7.1 [FJ3].

- (i) $\tau(1, 3, 3) = 5$,
- (ii) $\tau(2, 3, 3) = 10$,
- (iii) $\tau(k, 3, 3) = 4k + 3$ for any $k \geq 3$.

The theorem can be reformulated, as follows:

Theorem 7.1*. *Every 3-connected planar graph G of order at least $k \geq 3$ has a connected subgraph K of order k such that the degree in G of every vertex of K is at most $4k + 3$. The bound $4k + 3$ is best possible.* □

If the minimum degree of graphs of \mathcal{H} is increased to 4 then a slightly smaller bound is obtained by Fabrici.

Theorem 7.2 [Fa1]. $\tau(k, 4, 3) = 4k - 1$ for any $k \geq 4$. \square

In the paper [HW] of Hexel and Walther and that of Hexel [He] the reader can find several bounds on $\tau(k, \delta, \rho)$.

Here we provide a result that gives a necessary condition for a fixed family \mathcal{L} of connected plane graphs to be light in $\mathcal{P}(3, 3)$.

Theorem 7.3. *If \mathcal{L} is a finite family of connected plane graphs H such that $\Delta(H) \geq 3$ or $\delta(H) \geq 2$, then $\varphi(\mathcal{L}, \mathcal{P}(3, 3)) = +\infty$. That means that \mathcal{L} is not light in $\mathcal{P}(3, 3)$.*

Proof. Let K be the disjoint union of all graphs from the family \mathcal{L} . Let T_o be a plane triangulation of K , that is the graph obtained from K by inserting necessary edges into K to obtain a plane triangulation. The rest is the same as in the proof of Theorem 5.2, using K as H and m arbitrarily large. \square

Corollary 7.3.1. *If \mathcal{L} is a light family in $\mathcal{P}(3, 3)$ then \mathcal{L} contains a k -path for some k .* \square

Theorem 7.1 leads to the following problem:

Problem 7.2. *Find an optimal set \mathcal{S}_k of trees on k vertices such that $\varphi(\mathcal{S}_k, \mathcal{P}(3, 3)) = 4k + 3$.*

Applying Theorems 5.1 and 7.1 we can easily get the following.

Theorem 7.4. *If $\mathcal{L} = \{P_k, K_{1,3}\}$, then $\varphi(\mathcal{L}, \mathcal{P}(3, 3)) = 4k + 3$.* \square

The next theorem generalizes this result. For $i \geq 0$, let S_i denote a generalized 3-star with a central vertex of degree 3, where the three paths with common endpoint have $i + 1$ vertices. Obviously, $S_0 = K_1$ and $S_1 = K_{1,3}$.

Theorem 7.5 [JV11]. *Let k and i be integers with $k \geq 3$ and $1 \leq i \leq \frac{k}{2}$. If $\mathcal{L}_i = \{P_k, S_i\}$, then $\varphi(\mathcal{L}_i, \mathcal{P}(3, 3)) = \min\{5k, 4(k + i) - 1\}$.* \square

8. WEIGHT PROBLEMS

For a graph H contained in a graph in a family \mathcal{H} , one can also consider the smallest integer $w(H, \mathcal{H})$ having the property that every graph $G \in \mathcal{H}$ that contains a copy of H whose vertices have degrees in G that sum to at most $w(H, \mathcal{H})$. For a subgraph K of G , let $w_G(K) = \sum_{v \in V(K)} \deg_G(v)$; this is called the *weight* of K in G , (see e.g. [G3]).

Similarly $w(H, \mathcal{H})$ can be called the *weight of H in \mathcal{H}* , and we say that the graph H is light in \mathcal{H} if $w(H, \mathcal{H})$ is finite. Note that $w_G(K)$ and $\varphi_G(K)$ refer to the sum and the maximum over the same finite set. Thus $w(H, \mathcal{H})$ is finite if and only if $\varphi(H, \mathcal{H})$ is finite and the two definitions of a light graph are equivalent. Now we formulate

Problem 8.1. *Find the precise value of $w(H, \mathcal{H})$ for a given graph H and a family of graphs \mathcal{H} .*

The precise value $w(H, \mathcal{H})$ is known only for a few light graphs (and families). We start our survey with a beautiful recent result of Mohar [Mo1].

Theorem 8.1 [Mo1]. *Let \mathcal{P}_{ham} be the family of all planar hamiltonian graphs. Then*

$$w(P_k, \mathcal{P}_{\text{ham}}) = 6k - 1, k \geq 1.$$

Proof. For a planar hamiltonian graph H on n vertices, let C_n be a hamiltonian cycle through vertices v_1, v_2, \dots, v_n in order. Let R_i be the part of C_n on vertices $v_i, v_{i+1}, \dots, v_{i+k-1}$ (indices modulo n). Let $w(R_i) = \sum_{j=i}^{i+k-1} \deg_H(v_j)$ denote the weight of R_i in H (that is the sum of degrees in H of vertices of R_i). Then

$$\sum_{i=1}^n w(R_i) = k \sum_{v \in V(H)} \deg(v) = 2k|E(H)| \leq 2k(3n - 6).$$

The last inequality is the well known corollary (2.4) of Euler's formula. Hence, one of the paths, say R_j , has weight at most $2k(3n - 6)/n$. Since this is less than $6k$, we obtain $w(P_k, \mathcal{P}_{\text{ham}}) \leq 6k - 1$. On the other hand, there are 5-connected plane triangulations which contain precisely 12 vertices of degree 5, and all other vertices are 6-vertices. Moreover, the 5-vertices are as far away from each other as we like. This shows that $w(P_k, \mathcal{P}_{\text{ham}}) \geq 6k - 1$ and completes the proof. \square

Clearly 2-connected outerplanar graphs are hamiltonian. When using the idea of Mohar one can prove that every 2-connected outerplanar has a k -path of the weight at most $4k - 1$. The next theorem is by Fabrici [Fa2].

Theorem 8.2 [Fa2]. *Let $\mathcal{O}(2)$ be the family of all 2-connected outerplanar graphs. Then for every $k \geq 3$*

$$\varphi(P_k; \mathcal{O}(2)) = k + 3 \text{ and } w(P_k; \mathcal{O}(2)) = 4k - 2.$$

\square

Fabrici, Harant, and Jendroľ [FHJ] further developed the ideas of Mohar's proof to show that an upper bound that is linear in k also holds for a wider family of plane graphs.

Theorem 8.3 [FHJ]. *For $G \in \mathcal{P}(\delta, \rho)$ let $c(G)$ be the length of a longest cycle of G . Let k be an integer $3 \leq k \leq c(G)$. If $c(G) \geq \sigma|V(G)|$ for some positive number σ , then G contains a k -path P_k such that*

$$w_G(P_k) \leq \left(\left(\frac{2\rho}{\rho - 2} - \delta \right) \frac{1}{\sigma} + \delta \right) k - 1.$$

\square

For the families $\mathcal{P}(\delta, \rho)$, only trivial bounds on $w(\delta, \rho, H)$ are known except in a few cases. Trivially,

$$w(\delta, \rho; H) \leq |V(H)|\varphi(\delta, \rho; H)$$

where $w(\delta, \rho; H) = w(H; \mathcal{P}(\delta, \rho))$. This yields e.g. $w(5, 3; P_k) \leq w(4, 3; P_k) \leq 5k^2 - 7k$, for $k \geq 8$, see [FHJW]. Also $w(3, 4; P_k) \leq \frac{5}{2}k^2$, see [HJT] and $w(3, 5; P_k) \leq \frac{5}{3}k^2$, see [JO].

[Je3] Jendroľ formulated the following problem

Problem 8.2. *Determine the precise value of $w(3, 3; P_k)$.*

The precise values are known up to $k = 3$: $w(3, 3; P_1) = 5$, $w(3, 3; P_2) = 13$ (Kotzig [Ko1]), and $w(3, 3; P_3) = 21$ (Ando et al. [AIK]). The bounds on $w(3, \rho; P_k)$ for $3 \leq \rho \leq 5$ in the next theorem are due to Fabrici, Harant and Jendrof [FHJ]. The lower bound in case (i) is by Fabrici and Jendrof [FJ3].

Theorem 8.4 [FHJ]. *Let k be an integer, $k \geq 4$. Then*

- (i) $k \log_2 k \leq w(3, 3; P_k) \leq k^2 + 13k$.
- (ii) $\frac{1}{2}k \log_2 k + \mathcal{O}(k) \leq w(3, 4; P_k) \leq \frac{1}{2}(k^2 + 13k)$.
- (iii) $\frac{1}{8}k \log_2 k + \mathcal{O}(k) \leq w(3, 5; P_k) \leq \frac{1}{3}(k^2 + 13k)$. □

Let us notice that there is a contrast between hamiltonian and nonhamiltonian planar graphs. Mohar's Theorem 8.1 provides a sharp linear upper bound while constructions by Fabrici and Jendrof [FJ3] show that the exact value of $w(3, 3; P_k)$ is not linear. We believe that $w(\delta, \rho; P_k) = \mathcal{O}(k \log_2 k)$. Mohar [Mo] has constructed 3-connected planar graphs proving that $w(4, 3; P_k) \geq \frac{9}{16}k \log_2 k$ and $w(5, 3; P_k) \geq \frac{3}{10}k \log_2 k$.

Due to a result of Tutte [Tu] that every 4-connected planar graph is hamiltonian Theorem 8.1 has the following corollary

Theorem 8.5 [Mo1]. *Every n -vertex 4-connected planar graph contains a k -path P_k of weight $\leq 6k - 1$ for every $1 \leq k \leq n$. Moreover, this bound is tight.* □

Problem 8.2 can be formulated more generally

Problem 8.4. *Determine the precise value of $w(\delta, \rho; H)$ for all light graphs H in the family $\mathcal{P}(\delta, \rho)$ for all possible pairs (δ, ρ) .*

The known precise results concerning this problem not mentioned in Section 4 are listed here: Theorem 2.1 of Lebesgue yield: $w(3, 5; P_4) = 12$, and $w(3, 5; P_5) = 17$. Jendrof and Madaras [JM] proved $w(5, 3; P_4) = w(5, 3; K_{1,3}) = 23$, and recently Borodin and Woodall [BW1] showed that $w(5, \bar{3}; C_4) = 25$, and $w(5, \bar{3}; C_5) = 30$, and $w(5, 3; K_{1,4}) = 30$. Madaras [Ma2] proved that $w(4, \bar{3}; P_4) \leq 31$ and $w(5, \bar{3}; P_5) = 29$.

Analogously we can define the value $w(\mathcal{L}, \mathcal{H})$ for a finite family of connected graphs \mathcal{L} and a family \mathcal{H} having the properties mentioned in Section 7. Namely, every member of \mathcal{L} is a proper subgraph of at least one member of \mathcal{H} . Let $w(\mathcal{L}, \mathcal{H})$ be the smallest integer t with the property that every graph G in \mathcal{H} that has a subgraph belonging to \mathcal{L} has such a subgraph H whose vertices have degrees in G such that sum to at most $w(\mathcal{L}, \mathcal{H})$.

A nice result recently published by Enomoto and Ota can be read as follows: "Every 3-connected planar graph G of order at least k contains a connected subgraph H of order k such that the degree sum of vertices of H in G is at most $8k - 1$." More precisely, using our notation, they proved

Theorem 8.6 [EO]. *Let k be an integer, and \mathcal{T}_k be the family of all trees on k vertices. If $k \geq 4$, then*

$$8k - 5 \leq w(\mathcal{T}_k, \mathcal{P}(3, 3)) \leq 8k - 1$$

$$8k - 5 \leq w(\mathcal{T}_k, \mathcal{P}(4, 3)) \leq 8k - 3$$

$$7k - 4 \leq w(\mathcal{T}_k, \mathcal{P}(5, 3)) \leq 7k - 2. \quad \square$$

They expect the following to be true

Conjecture 8.5 [EO]. *If k is an integer and $k \geq 4$, then*

$$w(\mathcal{T}_k, \mathcal{P}(3, 3)) = 8k - 5. \quad \square$$

Madaras and Škrekovski [MŠ] investigated the conditions related to weight of a fixed subgraph of plane graphs that can enforce the existence of light graphs in families of plane graphs. For the families of plane graphs and triangulations whose edges are of weight at least w they study the necessary and sufficient conditions for lightness of certain graphs according to values of w . We like some of their results:

Theorem 8.7 [MŠ]. *Let $\mathcal{R}(w)$ be the family of all planar graphs of minimum degree at least 3 whose edges are of weight at least w .*

- (i) *The 4-path P_4 is light in $\mathcal{R}(w)$ if and only if $8 \leq w \leq 13$.*
- (ii) *The k -cycle C_k , $k \in \{3, 4\}$, is light in $\mathcal{R}(w)$ if and only if $10 \leq w \leq 13$.*
- (iii) *The star $K_{1,4}$ is light in $\mathcal{R}(w)$ if and only if $9 \leq w \leq 13$.* □

9. LIGHT SUBGRAPHS OF GRAPHS EMBEDDED ON SURFACES

In this section we discuss light subgraphs of connected graphs embedded into surfaces other than the plane. First we recall necessary terms.

Throughout this section we use terminology of [MT]. However, we recall some definitions. An *orientable surface* \mathbb{S}_g of genus g is obtained from the sphere by adding g handles. Correspondingly, a *nonorientable surface* \mathbb{N}_q of genus q is obtained from the sphere by adding q crosscaps. The Euler characteristic is defined by

$$\chi(\mathbb{S}_g) = 2 - 2g \text{ and } \chi(\mathbb{N}_q) = 2 - q.$$

By a *surface* \mathbb{M} we mean either an orientable surface \mathbb{S}_g or a nonorientable surface \mathbb{N}_q . By the *genus* g (the *nonorientable genus* q) of a graph G we mean the smallest integer g (q) such that G has an embedding into \mathbb{S}_g (\mathbb{N}_q , respectively).

If a graph G is embedded in a surface \mathbb{M} then the connected regions of $\mathbb{M} - G$ are called the *faces* of G . If each face is an open disc then the embedding is called a *2-cell embedding*. If each vertex has degree at least 3 and each vertex of degree h is incident with h different faces then G is called a *map* in \mathbb{M} . If, in addition, G is 3-connected and the embedding has "representativity" at least three, then G is called a *polyhedral map* in \mathbb{M} , see e.g. Robertson and Vitray [RV]. Let us recall that the *representativity* (or face width) of a (2-cell) embedded graph G into a surface \mathbb{M} is equal to the smallest number k such that \mathbb{M} contains a noncontractible closed curve that intersects the graph G in k points. We say that H is a *subgraph* of a polyhedral map G if H is a subgraph of the underlying graph of the map G .

For a map G let $V(G)$, $E(G)$ and $F(G)$ be the vertex set, the edge set and the face set G , respectively. For a map G on a surface \mathbb{M} Euler's famous formula states

$$|V(G)| - |E(G)| + |F(G)| = \chi(\mathbb{M}).$$

In 1990, Ivančo [Iv] generalized the Theorem 3.1 of Kotzig in the following way:

Theorem 9.1 [Iv]. *If G is a connected graph of orientable genus g and minimum degree at least 3, then G contains an edge e of weight*

$$w(e) \leq \begin{cases} 2g + 13 & \text{for } 0 \leq g \leq 3 \\ 4g + 7 & \text{for } g \geq 3. \end{cases}$$

Furthermore, if G does not contain 3-cycles, then

$$w(e) \leq \begin{cases} 8 & \text{for } g = 0 \\ 4g + 5 & \text{for } g \geq 1. \end{cases}$$

Moreover, all bounds are best possible. \square

An analogous result for graphs on non-orientable surfaces proved by JendroŔ and Tuhársky [JTU] is as follows

Theorem 9.2 [JTU]. *If G is a connected graph of minimum degree at least 3 on a non-orientable genus $q \geq 1$, then G contains an edge e of weight*

$$w(e) \leq \begin{cases} 2q + 11 & \text{for } 1 \leq q \leq 2, \\ 2q + 9 & \text{for } 3 \leq q \leq 5, \\ 2q + 7 & \text{for } q \geq 6. \end{cases}$$

Furthermore, if G does not contain 3-cycles, then

$$w(e) \leq 4q + 5 \quad \text{for } q \geq 1$$

Moreover, all bounds are best possible. \square

For the projective plane, the nonorientable surface of the smallest genus, using the same ideas as in the proof of Theorem 3.1, one can easily prove

Theorem 9.3. *Every connected projective planar graph of minimum degree at least 3 contains a $(3, a)$ -edge with $3 \leq a \leq 10$, or a $(4, b)$ -edge with $4 \leq b \leq 7$ or a $(5, c)$ -edge with $5 \leq c \leq 6$. The bounds 10, 7, and 6 are best possible. \square*

The bounds in Theorem 9.1 and 9.2 can be essentially improved if embedded graphs have a "large" number of vertices. Namely, the following holds:

Theorem 9.4 [JTV]. *Let G be a normal map on surface \mathbb{M} of Euler characteristic $\chi(\mathbb{M}) \leq 0$ and let n be the number of vertices of G . If*

- (a) $\sum(\deg_G(x) - 6) > 48|\chi(\mathbb{M})|$, or
- (b) $n > 26|\chi(\mathbb{M})|$,

then G contains an (a, b) -edge such that

- (i) $a = 3$ and $3 \leq b \leq 12$, or
- (ii) $a = 4$ and $4 \leq b \leq 8$, or
- (iii) $5 \leq a \leq 6$ and $5 \leq b \leq 6$.

The bounds 12, 8, and 6 are best possible. \square

Nothing seems to be done in the following

Problem 9.1. Find an analogue of Theorem 3.3 for polyhedral maps on manifolds \mathbb{M} of Euler characteristics $\chi(\mathbb{M}) \leq 0$.

For the projective plane, Sanders established sharp inequalities.

Theorem 9.5 [Sa]. Every normal projective planar map satisfies the following inequality:

$$\begin{aligned} 40e_{3,3} + 25e_{3,4} + 16e_{3,5} + 10e_{3,6} + \frac{20}{3}e_{3,7} + 5e_{3,8} + \frac{5}{2}e_{3,9} + 2e_{3,10} \\ + \frac{50}{3}e_{4,4} + 11e_{4,5} + 5e_{4,6} + \frac{5}{3}e_{4,7} \\ + \frac{16}{3}e_{5,5} + 2e_{5,6} \geq 60 \end{aligned}$$

□

and each of these coefficients is best possible.

In Theorem 9.3 we have the same coefficients as in the planar case (Theorem 3.2), but on the right side the value 120 has been weakened to 60.

In the subclass of all normal projective planar graphs with minimum degree at least 4 we have $e_{3,j} = 0$ for $3 \leq j \leq 10$. Sanders proved that in the resulting inequality all coefficients are best possible.

Theorem 9.6 [Sa]. Every normal projective plane map of minimum degree four satisfies the inequality

$$50e_{4,4} + 33e_{4,5} + 15e_{4,6} + 5e_{4,7} + 16e_{5,5} + 6e_{5,6} \geq 180,$$

and each of these coefficients is best possible. □

Theorem 9.7 [Sa]. Every projective planar graph of minimum degree five satisfies the inequality

$$16e_{5,5} + 7e_{5,6} \geq 210,$$

and each of these coefficients is best possible. □

If in Theorem 9.5 we use $e_{4,j} = 0$ for $4 \leq j \leq 7$ then an inequality is obtained which differs in the coefficient of $e_{5,5}$.

Theorem 9.8 [Sa]. Every projective plane graph of minimum degree five satisfies the inequality

$$18f_{5,5,5} + 9f_{5,5,6} + 5f_{5,5,7} + 4f_{5,6,6} \geq 72,$$

and each of these coefficients is best possible. □

Euler's formula implies (with some terms left out) $3v_3 + 2v_4 + v_5 \geq 12$ for the plane and $3v_3 + 2v_4 + v_5 \geq 6$ for the projective plane. Most of the above inequalities differ only on the right side, where 12 appears for the normal plane graphs and 6 for the normal projective planar graphs. This is completely true if the minimum degrees are 3 or 4, respectively. The only inequality that does not follow precisely these lines is the light edge inequality for graphs of minimum degree five. For the plane, the coefficient of $e_{5,5}$ went from 8/15

to $7/15$. For the projective plane, it is lowered from $8/15$ to $16/35$. Each other coefficient of the inequalities in the projective planar case is equal to the corresponding coefficient in the plane case.

Using the same arguments as for the planar case it is possible to prove the following analogues of Theorem 3.1 and Theorem 7.1

Theorem 9.9 [FJ3]. *Every 3-connected projective planar graph G that contains a k -path contains such a path whose vertices all have degree at most $5k$ in G . The bound $5k$ is best possible.* \square

Theorem 9.10 [JV5]. *For all $k \geq 3$ every 3-connected projective planar graph G of order at least k contains a connected subgraph H of order k whose vertices all have degree at most $4k + 3$ in G .* \square

We generalized these and other results on light subgraphs to surfaces \mathbb{M} with nonpositive Euler characteristic. For details on these results, see [JV10]. In the next subsections we give a brief survey only. We mention that all other theorems of Section 7 are also true for 3-connected projective planar graphs.

Theorem 9.11 [JV1]. *Each polyhedral map G on \mathbb{M} that contains a k -path contains such a k -path whose vertices all have degree at most $k \lfloor (5 + \sqrt{49 - 24\chi(\mathbb{M})})/2 \rfloor$ in G . Equality holds for even k .* \square

Let K_n and K_n^- denote the complete graph on n vertices and the graph obtained from it by deleting one edge, respectively. For odd k we can show:

Theorem 9.12 [JV4]. *For each odd k greater than $\frac{4}{3} \left\lfloor \frac{5 + \sqrt{49 - 24\chi(\mathbb{M})}}{2} \right\rfloor + 1$:*

- (i) *the upper bound in Theorem 9.11 is attained at infinite many orientable surfaces and at infinite nonorientable surfaces, where these surfaces are characterized by the fact that each of these surfaces has a triangular embedding of a K_n^- ;*
- (ii) *the upper bound in Theorem 9.11 is not attained at infinite many orientable surfaces and at infinite many nonorientable surfaces, where these surfaces are characterized by the fact that each of these surfaces has a triangular embedding of a K_n^- (in this case an upper bound is*

$$\left\lfloor \left(k - \frac{1}{3} \right) \frac{5 + \sqrt{49 - 24\chi(\mathbb{M})}}{2} \right\rfloor.$$

\square

A polyhedral map G is called *large* if it has a large number of vertices or large positive k -charge, where the *positive charge* of G is $\sum_{\deg_G(u) > 6k} (\deg_G(u) - 6k)$.

Theorem 9.13 [JV2], [JV3]. *Every large polyhedral maps on a surface \mathbb{M} of nonpositive Euler characteristic that contains a k -path contains such a k -path whose vertices all have*

degree, in G , at most $6k$ for $k = 1$ or even $k \geq 2$ and at most $6k - 2$ for odd $k \geq 3$. Moreover, these bounds are tight. \square

The upper bound on maximum degree of vertices of light paths polyhedral maps on a surface \mathbb{M} depends on $\sqrt{|\chi(\mathbb{M})|}$. In arbitrary embedding of 3-connected graphs (multigraphs) in \mathbb{M} this degree bound is a linear function of $|\chi(\mathbb{M})|$.

Theorem 9.14 [JV6]. *Each 3-connected multigraph G on \mathbb{M} without loops and 2-faces that has a k -path contains such a k -path whose vertices have degree at most $(6k - 2\varepsilon)(1 + |\chi(\mathbb{M})|/3)$ in G , where $\varepsilon = 0$ if $k \geq 2$ is even, and $\varepsilon = 1$ if $k \geq 3$ is odd. The bounds are best possible.* \square

Theorem 9.15 [JV7]. *For 3-connected graphs on \mathbb{M} the precise degree bound is*

$$2 + \lfloor (6k - 6 - 2\varepsilon)(1 + |\chi(\mathbb{M})|/3) \rfloor \text{ for } k \geq 4.$$

\square

Fabrici, Hexel, Jendroľ, and Walther [FHJW] proved that each 3-connected plane graph of minimum degree at least 4 that has k -path contains such a k -path whose vertices all have degree at most $5k - 7$ in G . This bound is sharp for $k \geq 8$. For surfaces other than the plane, we have

Theorem 9.16 [JV8]. *For $k \geq 8$, each large polyhedral map G on \mathbb{M} of minimum degree at least 4 that contains a k -path contains such a k -path whose vertices have degree at most $6k - 12$ in G . This bound is sharp for even k , and it must be at least $6k - 14$ for odd k .* \square

In 3-connected plane graphs of minimum degree at least 5 only the bound $5k - 7$ [FHJW] is known. For large polyhedral maps on 2-manifolds \mathbb{M} , the degree bound is not a linear function on k .

Theorem 9.17 [JV8]. *Let k be an integer at least 66. Each large polyhedral map G on \mathbb{M} of minimum degree at least 5 that contains a k -path contains such a k -path whose all vertices have degree at most $6k - \log_2 k + 2$. Moreover, the exact bound is at least $6k - 72 \log_2 k - 132$.* \square

In families of polyhedral maps of Theorem 9.10, 9.12 and 9.15 and in embeddings of 3-connected multigraphs (Theorem 9.13), and in embeddings of 3-connected graphs (Theorem 9.14) only k -paths are light for every k .

In the families of large polyhedral maps of minimum degree at least 5 on surfaces of nonpositive genus one can prove the existence of other light graphs. So the cycle C_3 is light there (see [JV12]), all proper spanning subgraphs H of the complete graph K_4 are light there, while K_4 itself is not light (see [JV13]). The 5-cycle C_5 and the 5-cycle with one or two diagonals are light in this class as well (see [JV14]). For other results see [JV9].

Fabrici and Jendroľ [FJ3] proved that each 3-connected plane graph G of order at least k contains a connected subgraph of order k whose vertices all have degree at most $4k + 3$. We have proved that this also holds for the projective plane. For polyhedral maps on \mathbb{M} , the degree bound for connected subgraphs of order k again depends on $\sqrt{|\chi(\mathbb{M})|}$. (This result is not presented here). For polyhedral maps with many vertices we proved

Theorem 9.18 [JV5]. *For $k \geq 2$, each polyhedral map on \mathbb{M} having at least $(8k^2 + 6k - 6)|\chi(\mathbb{M})| + 1$ vertices contains a connected subgraph of order k whose vertices all have degree at most $4k + 4$. This bound is best possible.* \square

For polyhedral maps the bound depends on $\sqrt{|\chi(\mathbb{M})|}$. In arbitrary embeddings of 3-connected graphs (multigraphs) in \mathbb{M} the bound is a linear function of $|\chi(\mathbb{M})|$.

Theorem 9.19 [JV6]. *For $k \geq 2$ each 3-connected multigraph G on \mathbb{M} that has no loops or 2-faces and has order at least k contains a connected subgraph of order k whose vertices all have degree at most $\lfloor (4k + 4)(1 + \frac{|\chi(\mathbb{M})|}{3}) \rfloor$ in G . This bound is sharp.* \square

Theorem 9.20 [JV7]. *For 3-connected graphs on \mathbb{M} , the precise degree bound is*

$$2 + \left\lfloor \left((4k - 2) \left(1 + \frac{|\chi(\mathbb{M})|}{3} \right) \right) \right\rfloor \text{ for } k \geq 5.$$

\square

Theorem 9.21 [JV6]. *For both large 3-connected multigraphs on \mathbb{M} without loops and 2-faces and for large 3-connected graphs on \mathbb{M} having at least $k \geq 2$ vertices the precise degree bound is $4k + 4$.* \square

We finish this section with an analogue of Theorem 8.6 recently proved by Kawarabayashi et al. [KNO].

Theorem 9.22 [KNO]. *For any non-spherical surface \mathbb{M} , any positive integer t there exist positive integers $r(\mathbb{M})$ and $n_0 = n_0(\mathbb{M}, t)$ such that if G is a n vertex, $n \geq n_0$, 3-connected graph embedded into \mathbb{M} with representativity $r(\mathbb{M})$, then G has a connected subgraph H of t vertices such that*

$$w_G(H) \leq \sum_{v \in V(H)} \deg_G(v) \leq 8t - 1.$$

10. RELATED TOPICS

The concept of the weight of an edge, of a face, of a path, or of a cycle as presented in this survey has served as a starting point for research in several other directions. We briefly mention some of them.

1. The idea of light edges was used by P. Erdős who formulated in 1990 at the conference in Prachatice (Czechoslovakia) the following max-min problem (see [IJ]): For a graph $G = (V, E)$ its edge weight $w(G)$ is defined as $\min\{w(e) | e \in E\}$. Let $\mathcal{G}(n, m)$ denote the family of all graphs with $n = |V|$ vertices and $m = |E|$ edges. Determine the value

$$W(n, m) = \max\{w(G) | G \in \mathcal{G}(n, m)\}.$$

Ivančo and JendroŔ [IJ] have proved some partial results. Recently JendroŔ and Schiermeyer [JS] have found a complete solution to Erdős's question and characterized all graphs on n vertices and m edges attaining this minimum weight.

A graph G from $\mathcal{G}(n, m)$ having no isolated vertices is *degree-constrained* if $a = \frac{2m}{n} < 2\delta$, where a is the average degree of G and $\delta = \delta(G)$ is minimum degree of G . Bose, Smid, and

Wood [BSW] proved that every degree-constrained graph has an edge uv with both $\deg(u)$ and $\deg(v)$ at most $\lfloor d \rfloor$ where $d = \frac{a\delta}{2\delta-a}$. Moreover they investigate matchings consisting of light edges in degree-constrained graphs.

2. The idea of a light path has been considered also in the family of all $K_{1,r}$ -free graphs, $r \geq 3$, that are graphs without $K_{1,r}$ as an induced subgraph. Harant et al. [HJRR] proved among others the following rather surprising result.

Theorem 10.1 [HJRR]. *If G is a $K_{1,r}$ -free graph on n vertices, where $r \geq 3$, then each induced path of length at least $2r - 1$ and each induced cycle of length at least $2r$ in G has the weight at most $(2r - 2)(n - \alpha_0)$, where α_0 is the independence number of G . Moreover, this bound is tight.* \square

3. For a polyhedral map G the p -vector is defined in [G1] to be a sequence $p = (f_i | i \geq 3)$ where f_i denotes the number of i -gonal faces of G . Rosenfeld [Ro] started to investigate the problem of characterization of p -vectors of 3-connected non-regular plane graphs (i.e. non-regular polyhedral graphs) whose edges have the same, constant, weight. Jendrol' and Jucovič [JJ] made first steps in dealing with this problem for polyhedral maps on orientable surfaces. There is a lot of open problems in this topic, see e.g. [Je1], [JJ].

4. Similarly, Jucovič [IT] suggested studying polyhedral maps with constant weight of faces. By the weight of a face α we mean the sum of degrees of vertices incident with α . All Platonic solids and all duals to Archimedean solids have constant face weights. Ivančo and Trenkler [IT], and Horňák and Ivančo [HI] determined the number of nonisomorphic 3-connected plane graphs having prescribed face weight w . There are infinitely many such graphs if and only if $16 \leq w \leq 21$ or $w = 23$ or $w = 25$. There is exactly one such graph if $w = 9$ or $w = 11$, for $w = 14$ there are four, and for $w = 15$ there are ten such graphs. For other values of w at least 12 there are exactly two such graphs. For $w = 10$ no such graph exists. Nothing is known about polyhedral maps with constant face weight on surfaces other than the plane.

5. The idea of the weight of an edge $e = uv$ being the degree sum of the endvertices u and v motivated Jendrol' and Ryjáček [JR] to introduce the concept of *tolerance* of the edge e . The *tolerance* $\tau(e)$ of the edge $e = uv$ is defined to be the absolute value of the difference of degrees of the vertices u and v ,

$$\tau(e) = |\deg(u) - \deg(v)|.$$

Necessary and sufficient conditions for the existence of connected planar and 3-connected planar graphs with constant edge tolerance appear in [JR]. Several constructions of graphs with constant edge tolerance for general graphs appear in Acharya and Vartak [AV]. An open problem is to find necessary and sufficient conditions for graphs with constant edge tolerance embedded into surfaces different from the sphere.

6. Motivated by "light" results, Mohar [Mo2] considered such problems for infinite planar graphs. He uses the discharging method to prove some new results in this direction. The general outline of the method is presented in [Mo2]. Many applications are given there, including results on light subgraphs and the following: Planar graphs with only finitely

many vertices of degree at most 5 and with subexponential growth contain arbitrarily large finite submaps of the tessellation of the plane or of some tessellation of the cylinder by equilateral triangles.

7. We feel a need to write down few remarks concerning the Lebesgue theorem (Theorem 2.1). It was published in 1940 but it remained unnoticed until 1967 when Ore's book [Or] appeared. Ore was aware of the importance of the theorem and therefore he included it into his book together with a complete proof (using Lebesgue's theory of the Euler's contributions) and with corrolaries. But only after its application in an Ore and Plummer's [OP] problem on cyclic coloring of plane graphs by Plummer and Toft [PT] the theorem started to attract people. (Note that the *cyclic colouring* is a colouring of vertices of plane graphs in such a way that the vertices incident with the same face receive different colours.) To prove the conjecture of Plummer and Toft [PT], see also [JT], that every 3-connected plane graph G has a cyclic colouring of its vertices with $\Delta^* + 2$ colours, there were attempts to improve some terms in the theorem. Here Δ^* means the size of the largest face in G .

Recall that the unavoidable set by Lebesgue's theorem (see Theorem 2.1) consists of six infinite series of faces, namely series of $(3, b, c)$ -triangles for $3 \leq b \leq 6$ and $c \geq 3$, $(4, 4, c)$ -triangles for $c \geq 4$, and $(3, 3, 3, d)$ -quadrangles for $d \geq 3$, and 126 individual faces. Horňák and Jendrol' [HJ1] reduced two infinite series of $(3, b, c)$ -triangles for $5 \leq b \leq 6$ and $c \geq 5$ to finite ones. Namely they proved the existence of an unavoidable set of configurations consisting of four infinite series and 160 individual configurations.

Note that, in general, none of infinite series of $(3, b, c)$ -triangles for $3 \leq b \leq 4, c \geq 3$, $(4, 4, c)$ -triangles for $c \geq 4$, and $(3, 3, 3, d)$ -quadrangles for $k \geq 3$ can be omitted. In [HJ2] Horňák and Jendrol' replaced the serie of $(4, 4, c)$ -triangles with few individual terms together with a configuration consisting of a chain of three quadrangles.

On the other side Borodin [Bo14] succesfully reduced individual terms of the theorem to 95 by letting all six infinite series $(3, b, c)$ -triangles for $3 \leq b \leq 6$, $(4, 4, c)$ -triangles, and $(3, 3, 3, c)$ -quadrangles, all with $c \geq 3$. In his paper [Bo14] Borodin posed the following problem: Find the best possible version(s) of Lebesgue's theorem.

For other discussions concerning the Lebesgue theorems readers are recommended to [Bo14], [BoL], and [HJ1]. (For a present situation concerning the conjecture of Plummer and Toft, see e.g. [EHJ].)

Acknowledgements.

Support of Slovak VEGA Grant 1/0424/03 is acknowledged. The authors wish to thank the anonymous referee for his careful reading of the paper and very detailed proposals that helped to improve the presentation of the paper.

REFERENCES

- [AV] B. D. Acharya, M. N. Vartak, *On the construction of graphs with given constant valence-difference (S) on each of their lines*, *Wiss. Z. TH Ilmenau* **26** No. 6 (1977), 33–60.
- [ABG] B. A. Aksenov, O. V. Borodin, A. N. Glebov, *On a structural property of plane graphs*, *Diskretn. Anal. Issled. Oper. Ser. 1*, **7** (2000), 5–16, (Russian).
- [AIK] K. Ando, S. Iwasaki, A. Kaneko, *Every 3-connected planar graph has a connected subgraph with small degree sum*, *Annual Meeting in Mathematical Society of Japan* (1993), (Japanesse).

- [AH] K. Appel, W. Haken, *Every planar map is four colorable*, Contemp. Math. **98**, Providence, RI: American Mathematical Society (AMS). XV, (1989), 751.
- [BLW] N. Z. Biggs, E. K. Lloyd, R. J. Wilson, *Graph Theory 1736-1936*, Clarendon Press, Oxford (1998).
- [Bo1] O. V. Borodin, *Solution of problems of Kotzig and Grünbaum concerning the isolation of cycles in planar graphs*, Math. Notes **46** (1989), 9–12, (Russian).
- [Bo2] O. V. Borodin, *On the total coloring of planar graphs*, J. Reine Angew. Math. **394** (1989), 180–185.
- [Bo3] O. V. Borodin, *Computing light edges in planar graphs*, in Topics in Combinatorics and Graph Theory (1990), 137–144, (R. Bodendieck and R. Henn Eds.) Physica-Verlag, Heidelberg.
- [Bo4] O. V. Borodin, *Structural properties and colorings of plane graphs*, Ann. Discrete Math. **51** (1992), 31–37.
- [Bo5] O. V. Borodin, *Precise lower bound for the number of edges of minor weight in planar maps*, Math. Slovaca **42** (1992), 129–142.
- [Bo6] O. V. Borodin, *Structural properties of planar maps with minimum degree 5*, Math. Nachr. **158** (1992), 109–117.
- [Bo7] O. V. Borodin, *Joint extension of two theorems of Kotzig on 3-polytopes*, Combinatorica **13** (1993), 121–125.
- [Bo8] O. V. Borodin, *Simultaneous coloring of edges and faces of plane graphs*, Discrete Math. **128** (1994), 21–33.
- [Bo9] O. V. Borodin, *Triangles with restricted degree sum of their boundary vertices in plane graphs*, Discrete Math. **137** (1995), 45–51.
- [Bo10] O. V. Borodin, *More about the weight of edges in planar graphs*, Tatra Mt. Math. Publ. **9** (1996), 11–14.
- [Bo11] O. V. Borodin, *Minimal vertex degree sum of a 3-path in plane maps*, Discuss. Math. Graph Theory **17** (1997), 279–284.
- [Bo12] O. V. Borodin, *Triangulated 3-polytopes without faces of low weight*, Discrete Math. **186** (1998), 281–285.
- [Bo13] O. V. Borodin, *Colourings and topological representations of graphs*, Diskretn. Anal. Issled. Oper. **3** (1996), 3–27, (Russian).
- [Bo14] O. V. Borodin, *An improvement of Lebesgue’s theorem on the structure of minor faces of 3-polytopes*, Diskretn. Anal. Issled. Oper. **9**, No. 3 (2002), 29–35, (Russian).
- [BoL] O. V. Borodin, D. V. Loparev, *The height of small faces in normal plane maps*, Diskretn. Anal. Issled. Oper. **5** (1998), 6–17, (Russian).
- [BS] O. V. Borodin, D. P. Sanders, *On light edges and triangles in planar graphs of minimum degree five*, Math. Nachr. **170** (1994), 19–24.
- [BW1] O. V. Borodin, D. R. Woodall, *Short cycles of low weight in normal plane maps with minimum degree five*, Discuss. Math. Graph Theory **18** (1998), 159–164.
- [BW2] O. V. Borodin, D. R. Woodall, *Cyclic degrees of 3-polytopes*, Graphs Combin. **15** (1999), 267–277.
- [BB] O. V. Borodin, H. Broersma, A. N. Glebov, J. van den Heuvel, *The structure of plane triangulations in terms of clusters and stars*, Diskretn. Anal. Issled. Oper. Ser. 1. **8** (2001), 15–39, (Russian).
- [BSW] P. Bose, M. Smid, D. R. Wood, *Light edges in degree-constrained graph*, Discrete Math., (submitted 2002).
- [EHJ] H. Enomoto, M. Horňák, S. Jendroľ, *Cyclic chromatic number of 3-connected plane graphs*, SIAM J. Discrete Math. Vol. 14 No. 1 (2001), 121–137.
- [EO] H. Enomoto, K. Ota, *Connected subgraphs with small degree sum in 3-connected planar graphs*, J. Graph Theory **30** (1999), 191–203.
- [Fa1] I. Fabrici, *On Vertex-Degree Restricted Subgraphs in Polyhedral Graphs*, Discrete Math. **256** (2002), 105–114.
- [Fa2] I. Fabrici, *Light graphs in families of outerplanar graphs*, (submitted).
- [FJ1] I. Fabrici, S. Jendroľ, *An inequality concerning edges of minor weight in convex 3-polytopes*, Discuss. Math. Graph Theory **16** (1996), 81–87.

- [FJ2] I. Fabrici, S. JendroŤ, *Subgraphs with restricted degrees of their vertices in planar 3-connected graphs*, Graphs Combin. **13** (1997), 245–250.
- [FJ3] I. Fabrici, S. JendroŤ, *Subgraphs with restricted degrees of their vertices in planar graphs*, Discrete Math. **191** (1998), 83–90.
- [FHJ] I. Fabrici, J. Harant, S. JendroŤ, *Paths of low weight in planar graph*, (submitted).
- [FHJW] I. Fabrici, E. Hexel, S. JendroŤ, H. Walther, *On vertex-degree restricted paths in polyhedral graphs*, Discrete Math. **212** (2000), 61–73.
- [Fr] P. Franklin, *The four color problem*, Amer. J. Math. **44** (1922), 225–236.
- [G1] B. Grünbaum, *Convex polytopes*, Interscience, New York (1967) or Springer, New York (2003),, Second Edition.
- [G2] B. Grünbaum, *Acyclic colorings of planar graphs*, Israel J. Math. **14** (1973), 390–408.
- [G3] B. Grünbaum, *Polytopal graphs*, in Studies in Graph Theory (D. R. Fulkerson, ed.), MAA Studies in Mathematics **12** (1975), 201–224.
- [G4] B. Grünbaum, *New views on some old questions of combinatorial geometry*, Int. Teorie Combinatorie, Rome, **1** (1976), 451–468.
- [GS] B. Grünbaum, G. C. Shephard, *Analogues for tiling of Kotzig’s theorem on minimal weights of edges*, Ann. Discrete Math. **12** (1982), 129–140.
- [HK] A. Hackmann, A. Kemnitz, *List edge colorings of outerplanar graphs*, Ars Combin. **60** (2001), 181–185.
- [HJT] J. Harant, S. JendroŤ, M. Tkáč, *On 3-connected plane graphs without triangular faces*, J. Combin. Theory Ser. B **77** (1999), 150–161.
- [HJRR] J. Harant, S. JendroŤ, B. Randerath, Z. Ryjáček, I. Schiermeyer, M. Voigt, *On weight of induced paths and cycles in claw-free graphs*, J. Graph Theory **36** (2001), 131–143.
- [H1] H. Heesch, *Untersuchungen zum Vierfarbenproblem*, Hochschulsriptum 810/a/b, Bibliographisches Institut, Mannheim 1969.
- [HM] J. van den Heuvel, S. McGuinness, *Coloring the square of a planar graph*, J. Graph Theory **42** (2003), 110–124.
- [He] E. Hexel, *On light in the family of 4-connected planar graphs*, Discrete Math. **251** (2002), 103–107.
- [HS] E. Hexel, R. Soták, *On light cycles in the family of 5-connected triangulation*, Manuscript, June 2003.
- [HW] E. Hexel, H. Walther, *On vertex-degree restricted paths in 4-connected planar graphs*, Tatra Mt. Math. Publ. **18** (1999), 1–13.
- [HI] M. Horňák, J. Ivančo, *On the number of 3-polytopes with constant face weight*, Studia Sci Math. Hungar. **39** (2002), 1–19.
- [HJ1] M. Horňák, S. JendroŤ, *Unavoidable set of face types for planar maps*, Discuss. Math. Graph Theory **16** (1996), 123–141.
- [HJ2] M. Horňák, S. JendroŤ, *On a Conjecture by Plummer and Toft*, Inc. J. Graph Theory **30** (1999), 177–189.
- [Iv] J. Ivančo, *The weight of a graph*, Ann. Discrete Math. **51** (1992), 113–116.
- [IJ] J. Ivančo, S. JendroŤ, *On extremal problems concerning weights of edges of graphs*, Colloq. Math. Soc. János Bolyai **60** Sets, Graphs and Numbers, Budapest (1992), 339–410.
- [IT] J. Ivančo, M. Trenkler, *3-polytopes with constant face weight*, Studia Sci. Math. Hungar. **35** (1999), 1–15.
- [Je1] S. JendroŤ, *Convex 3-polytopes with constant edge weight*, Colloq. Math. Soc. János Bolyai **63** Intuitive geometry, Szeged (Hungary, 1991) North Holland, Budapest (1994), 173–179.
- [Je2] S. JendroŤ, *A structural property of convex 3-polytopes*, Geom. Dedicata **68** (1997), 91–99.
- [Je3] S. JendroŤ, *Light paths in 3-connected planar graphs (Problem 294)*, Discrete Math. **191** (1998), 248.
- [Je4] S. JendroŤ, *A short proof of Kotzig’s theorem on minimal edge weights of convex 3-polytopes*, Proc. Internat. Scient. Conference of Math. (V. Balint Ed.) Žilina (1999), 35–38.
- [Je5] S. JendroŤ, *Triangles with restricted degrees of their boundary vertices in plane triangulations*, Discrete Math. **196** (1999), 177–196.

- [Je6] S. Jendroř, *Paths with restricted degrees of their vertices in planar graphs*, Czechoslovak Math. J. **49** (1999), 481–490, (124).
- [JJ] S. Jendroř, E. Jucovič, *On face-vectors of maps with constant weight of edges*, Studia Sci. Math. Hungar. **17** (1982), 159–175.
- [JM] S. Jendroř, T. Madaras, *On light subgraphs in plane graphs of minimum degree five*, Discuss. Math. Graph Theory **16** (1996), 207–217.
- [JO] S. Jendroř, P. Owens, *On light graphs in 3-connected plane graphs without triangular or quadrangular faces*, Graphs Combin. **17** (2001), 659–680.
- [JR] S. Jendroř, Z. Ryjáček, *3-polytopes of constant tolerance of edges*, Comment. Math. Univ. Carolin. **22,4** (1981), 843–850.
- [JS] S. Jendroř, I. Schiermeyer, *On max-min problem concerning weights of edges*, Combinatorica **21** (2001), 351–359.
- [JT_u] S. Jendroř, M. Tuhářsky, *A Kotzig type theorem for non-orientable surfaces*, (submitted).
- [JV1] S. Jendroř, H.-J. Voss, *A local property of polyhedral maps on compact 2-dimensional manifolds*, Discrete Math. **212** (2000), 111–120.
- [JV2] S. Jendroř, H.-J. Voss, *A local property of large polyhedral maps on compact 2-dimensional manifolds*, Graphs Combin. **15** (1999), 303–313.
- [JV3] S. Jendroř, H.-J. Voss, *Light paths with an odd number of vertices in large polyhedral maps*, Ann. Comb. **2** (1998), 313–324.
- [JV4] S. Jendroř, H.-J. Voss, *Light paths with an odd number of vertices in polyhedral maps*, Czechoslovak Math. J. **50(120)** (2000), 555–564.
- [JV5] S. Jendroř, H.-J. Voss, *Subgraphs with restricted degrees of their vertices in large polyhedral maps on compact 2-manifolds*, European J. Combin. **20** (1999), 821–832.
- [JV6] S. Jendroř, H.-J. Voss, *Light subgraphs of multigraphs on compact 2-dimensional manifolds*, Discrete Math. **233** (2001), 329–351.
- [JV7] S. Jendroř, H.-J. Voss, *Two local and one global properties of 3-connected graphs on compact 2-dimensional manifolds*, J. Combin. Theory Ser. B, (submitted).
- [JV8] S. Jendroř, H.-J. Voss, *Light paths in large polyhedral maps with prescribed minimum degree*, Australas. J. Combin. **25** (2002), 79–102.
- [JV9] S. Jendroř, H.-J. Voss, *Subgraph with restricted degrees of their vertices in polyhedral maps on compact 2-manifolds*, Ann. Comb. **5** (2001), 211–226.
- [JV10] S. Jendroř, H.-J. Voss, *Light subgraphs of graphs embedded in 2-dimensional manifolds of Euler characteristic ≤ 0 - a survey*, In "Paul Erdős and his Mathematics, I" (G. Halász, L. Lovász, M. Simonovits, T. Sos, Eds.), Springer, Budapest (Hungary) 1999, Bolyai Society Mathematicae Studies, 11. Budapest (2002), 375–411.
- [JV11] S. Jendroř, H.-J. Voss, *Light classes of generalized stars in polyhedral maps on surfaces*, Discuss. Math. Graph Theory **24** (2004), 85–107.
- [JV12] S. Jendroř, H.-J. Voss, *Light subgraphs of order ≤ 3 in large maps of minimum degree 5 on compact 2-manifold*, European J. Combin., (to appear).
- [JV13] S. Jendroř, H.-J. Voss, *Light subgraphs of order 4 in large maps of minimum degree 5 on compact 2-manifold*, Australas. J. Combin. **28** (2003), 171–199.
- [JV14] S. Jendroř, H.-J. Voss, *The 5-cycle C_5 is light in large maps of minimum degree 5 on compact 2-manifolds*, Bul. Acad. Stiinte Repub. Mold. Mat. **3(40)** (2002), 106–124.
- [JTV] S. Jendroř, M. Tuhářsky, H.-J. Voss, *A Kotzig type theorem for large maps on surfaces*, Tatra Mt. Math. Publ. **27** (2003), 153–162.
- [JMST] S. Jendroř, T. Madaras, R. Soták, Z. Tuza, *On light cycles in plane triangulations*, Discrete Math. **197/198** (1999), 453–467.
- [JT] T. R. Jensen, B. Toft, *Graph coloring problems*, John-Wiley Sons, New York (1995).
- [Ju1] E. Jucovič, *Strengthening of a theorem about 3-polytopes*, Geom. Dedicata **13** (1974), 233–237.
- [Ju2] E. Jucovič, *Convex polytopes*, Veda Bratislava, 1981, (Slovak).
- [KNO] K. Kawarabayashi, A. Nakamoto, K. Ota, *Subgraphs of graphs on surfaces with high representativity*, J. Combin. Theory Ser. B **89** (2003), 207–229.

- [Ko1] A. Kotzig, *Contribution to the theory of Eulerian polyhedra*, Mat. Čas SAV (Math. Slovaca) **5** (1955), 111–113, (Slovak).
- [Ko2] A. Kotzig, *From the theory of eulerian polyhedrons*, Mat. Čas SAV (Math. Slovaca) **13** (1963), 20–34, (Russian).
- [Ko3] A. Kotzig, *Extremal polyhedral graphs*, Ann. New York Acad. Sci. **319** (1979), 565–570.
- [Le] H. Lebesgue, *Quelques conséquences simples de la formule d'Euler*, J. Math. Pures Appl. **19** (1940), 27–43.
- [Ma1] T. Madaras, *Note on weights of paths in polyhedral graphs*, Discrete Math. **203** (1999), 267–269.
- [Ma2] T. Madaras, *Note on the weight of paths in plane triangulation of minimum degree 4 and 5*, Discuss. Math. Graph Theory **20** (2000), 173–180.
- [MŠ] T. Madaras, R. Škrekovski, *Heavy paths, light stars, and big melons*, Discrete Math., (to appear).
- [MaŠV] B. Madaras, R. Škrekovski, H.-J. Voss, *The 7-cycle C_7 is light in the family of planar graphs with minimum degree 5*, Discrete Math., (to appear).
- [Mk] J. Malkevitch, *Polytopal graphs*, Selected topics in Graph theory **3** (L.W. Beineke and R.J. Wilson, Eds.), Academic Press London, (1988), 169–188.
- [Mo1] B. Mohar, *Light paths in 4-connected graphs in the plane and other surfaces*, J. Graph Theory **34** (2000), 170–179.
- [Mo2] B. Mohar, *Light structures in infinite planar graphs without the strong isoperimetric property*, Trans. Amer. Math. Soc. **354** (2002), 3059–3074.
- [MT] B. Mohar, C. Thomassen, *Graphs on Surfaces*, The Johns Hopkins University Press, Baltimore and London (2001).
- [MŠV] B. Mohar, R. Škrekovski, H.-J. Voss, *Light subgraphs in planar graphs of minimum degree 4 and edge-degree 9*, J. Graph Theory **44** (2003), 261–295.
- [MS] M. Molloy, M. R. Salavatipour, *A bound on the chromatic number of the square of a planar graph*, Manuscript 2003.
- [Or] O. Ore, *The four color problem*, Academic Press, New York/London (1967).
- [OP] O. Ore, M. D. Plummer, *Cyclic coloration of plane graphs*, in Recent Progress in Combinatorics, W.T. Tutte, ed., Academic Press, New York (1969), 287–293.
- [Pl] M. D. Plummer, *On the cyclic connectivity of planar graph* (1972), Graph Theory and Application, Springer, Berlin, 235–242.
- [PT] M. D. Plummer, B. Toft, *Cyclic coloration of 3-polytopes*, J. Graph Theory **11** (1987), 507–515.
- [RV] N. Robertson, R. P. Vitray, *Representativity of surface embeddings*, in "Paths, Flows and VLSI - Layout" (B. Korte, L. Lovász, J. Prömel, and A. Schrijver, Eds.) Springer-Verlag, Berlin/New York (1990), 293–328.
- [RSST] N. Robertson, D. Sanders, P. Seymour, R. Thomas, *The four-color theorem*, J. Combin. Theory Ser. B **70** (1997), 2–44.
- [Ro] M. Rosenfeld, *Polytopes of constant weight*, Israel J. Math. **21** (1975), 24–30.
- [Sa] D. P. Sanders, *On light edges and triangles in projective planar graphs*, J. Graph Theory **21** (1996), 335–342.
- [SR] E. Steinitz, H. Rademacher, *Vorlesung über die Theorie der Polyeder*, Berlin (1934).
- [Tu] W. T. Tutte, *A theorem on planar graphs*, Trans. Amer. Math. Soc. **82** (1956), 99–116.
- [Wg] G. Wegner, *Graphs with given diameter and a coloring problem*, Technical report, University of Dortmund (1977).
- [WL1] W. Weifan, K.-W. Lih, *Structural properties and edge choosability of planar graphs without 6-cycles*, Combinatorics, Probability and Computing **10** (2001), 267–276.
- [We] P. Wernicke, *Über den kartographischen Vierfarbensatz.*, Math. Ann. **58** (1904), 413–426.
- [WW] D. R. Woodall, R. J. Wilson, *The Appel-Haken proof of the Four-Color Theorem*, in "Selected Topics in Graph Theory", (L.W. Beineke and R.J. Wilson, Eds.) Academic Press London-New York-San Francisco (1978), 83–101.
- [Za1] J. Zaks, *Extending Kotzig's Theorem*, Israel J. Math. **45** (1983), 281–296.
- [Za2] J. Zaks, *Extending two theorems of A. Kotzig*, Discrete Math. **43** (1983), 309–315.
- [Zi] G. M. Ziegler, *Lectures on polytopes* (1995), Grad. Texts in Math. **152**, Springer-Verlag, Berlin.